

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 September 2002 (06.09.2002)

PCT

(10) International Publication Number
WO 02/068646 A2

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/54, 14/715, G01N 33/68, C07K 16/28, 16/24

(74) Agent: **GRIPP, Sandy, S.**; Schering Plough Corporation,
Patent Department- K-6-1- 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(21) International Application Number: **PCT/US01/50351**

(22) International Filing Date:
9 November 2001 (09.11.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/247,218 10 November 2000 (10.11.2000) US
60/298,268 14 June 2001 (14.06.2001) US

(71) Applicant: **SCHERING CORPORATION** [US/US];
Patent Department - K-6-1 1990, 2000 Galloping Hill
Road, Kenilworth, NJ 07033-0530 (US).

(72) Inventors: **RECHE-GALLARDO, Pedro, A.**; 96 Linden
Street, # 6, Allston, MA 02134 (US). **DE WAAL MALE-
FYT, Rene**; 891 Muender Avenue, Sunnyvale, CA 94086
(US). **BAZAN, J., Fernando**; 426 Waverly Street, #6, Palo
Alto, CA 94301 (US). **KASTELEIN, Robert, A.**; 463
Summit Drive, Redwood City, CA 94062 (US). **LIU, Yong-
Jun**; 4010 Villa Vista, Palo Alto, CA 94306 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PH, PL, PT,
RO, RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ,
VN, YU, ZA.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).

Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

WO 02/068646 A2

(54) Title: **MAMMALIAN CYTOKINES; RECEPTORS; RELATED REAGENTS AND METHODS**

(57) Abstract: Nucleic acids encoding mammalian cytokine receptor, e.g. for cytokine IL-B50, purified proteins and fragments thereof. Antibodies, both polyclonal and monoclonal, are also provided. Methods of using the compositions for both diagnostic and therapeutic utilities are described.

MAMMALIAN CYTOKINES; RECEPTORS; RELATED REAGENTS AND METHODS

FIELD OF THE INVENTION

The present invention relates to compositions and methods for affecting mammalian
5 physiology, including hematopoietic cell proliferation or immune system function. In
particular, it provides methods of using nucleic acids, proteins, and antibodies which
regulate development and/or the immune system; and provides functional details on ligand-
receptor pairing. Diagnostic and therapeutic uses of these materials are also disclosed.

BACKGROUND OF THE INVENTION

Recombinant DNA technology refers generally to techniques of integrating genetic
information from a donor source into vectors for subsequent processing, such as through
introduction into a host, whereby the transferred genetic information is copied and/or
expressed in the new environment. Commonly, the genetic information exists in the form of
15 complementary DNA (cDNA) derived from messenger RNA (mRNA) coding for a desired
protein product. The carrier is frequently a plasmid having the capacity to incorporate
cDNA for later replication in a host and, in some cases, actually to control expression of the
cDNA and thereby direct synthesis of the encoded product in the host.

For some time, it has been known that the mammalian immune response is based on
20 a series of complex cellular interactions, called the "immune network". Recent research has
provided new insights into the inner workings of this network. While it remains clear that
much of the immune response does, in fact, revolve around the network-like interactions of
lymphocytes, macrophages, granulocytes, and other cells, immunologists now generally
hold the opinion that soluble proteins, known as lymphokines, cytokines, or monokines,
25 play critical roles in controlling these cellular interactions. Thus, there is considerable
interest in the isolation, characterization, and mechanisms of action of cell modulatory
factors, an understanding of which will lead to significant advancements in the diagnosis
and therapy of numerous medical abnormalities, e.g., immune system disorders.

Lymphokines apparently mediate cellular activities in a variety of ways. They have
30 been shown to support the proliferation, growth, and/or differentiation of pluripotent
hematopoietic stem cells into vast numbers of progenitors comprising diverse cellular
lineages which make up a complex immune system. Proper and balanced interactions

between the cellular components are necessary for a healthy immune response. The different cellular lineages often respond in a different manner when lymphokines are administered in conjunction with other agents.

Cell lineages especially important to the immune response include two classes of lymphocytes: B-cells, which can produce and secrete immunoglobulins (proteins with the capability of recognizing and binding to foreign matter to effect its removal), and T-cells of various subsets that secrete lymphokines and induce or suppress the B-cells and various other cells (including other T-cells) making up the immune network. These lymphocytes interact with many other cell types. Monocytes are precursors of macrophages which, with dendritic cells, are functionally important in their roles as processors and presenters of antigen, an important step in initiation of an immune response.

IL-7 is a cell modulatory factor which affects hematopoietic cell growth and/or differentiation. See, e.g., Mire-Sluis and Thorpe (1998) Cytokines Academic Press, San Diego; Thomson (ed. 1998) The Cytokine Handbook (3d ed.) Academic Press, San Diego; Metcalf and Nicola (1995) The Hematopoietic Colony Stimulating Factors Cambridge University Press; and Aggarwal and Gutterman (1991) Human Cytokines Blackwell.

Research to better understand and treat various immune disorders has been hampered by the general inability to maintain cells of the immune system in vitro. Immunologists have discovered that culturing many of these cells can be accomplished through the use of T-cell and other cell supernatants, which contain various growth factors, including many of the lymphokines.

From the foregoing, it is evident that the understanding of the signal transduction pathways and identification of components in such pathways should contribute to new therapies for a wide range of degenerative or abnormal conditions which directly or indirectly involve development, differentiation, or function, e.g., of the immune system and/or hematopoietic cells. Furthermore, soluble regulatory molecules, including cytokines, are known to sometimes act outside the immune system with effects on physiology (leptin), morphogenesis, and tissue and skeletal remodeling (RANKL). Thus, the discovery and understanding of novel cytokine-like molecules and their receptors which enhance or potentiate the beneficial activities of other lymphokines would be highly advantageous.

SUMMARY OF THE INVENTION

The present invention is based, in part, on the discovery of the receptor complex for the cytokine ligand IL-B50. Moreover, identification of the components allows for the identification of cell types and stages which express the necessary components to be responsive to ligand. This provides insights and capacity to predict the physiological and immunological role of the cytokine.

The present invention provides, e.g., methods of producing a ligand:receptor complex, comprising contacting: a substantially pure or recombinant mammalian IL-B50 with a receptor comprising the IL-7R α or the R δ 2 subunit; a mammalian IL-B50 with a receptor comprising a substantially pure or recombinant IL-7R α subunit; or a mammalian IL-B50 with a receptor comprising a substantially pure or recombinant R δ 2 subunit; which contacting thereby allows the complex to form. In preferred embodiments, the mammalian IL-B50 is primate IL-B50, such as human IL-B50; the complex formation results in signal transduction, STAT activation, or TARC expression; the receptor is on a cell; the receptor comprises both IL-7R α and R δ 2 subunit; the complex formation results in a physiological change in the cell expressing the receptor; the contacting is in combination with a proliferative agent, cytokine, or chemokine; the contacting allows quantitative detection of the ligand; or receptor is on a hematopoietic cell, including a lymphoid lineage cell, a myeloid cell such as a monocyte, or dendritic cell.

Another method is provided for modulating physiology or development of an IL-7R α or R δ 2 expressing cell comprising contacting the cell to an exogenous agonist or antagonist of a mammalian IL-B50. Various embodiments include those wherein: the antagonist is an antibody which neutralizes the mammalian IL-B50, a mutein of the IL-B50; or an antibody which binds to IL-7R α or R δ 2 or a complex of both; or the physiology is selected from proliferation, lymphoid lineage cell development, antigen presentation, or production of inflammatory mediators, including cytokines, chemokines, or adhesion molecules; or the cell is a hematopoietic cell. Other embodiments include those wherein: the antagonist is an antibody and the physiology is hematopoietic cell proliferation; the agonist is IL-B50 and the physiology is hematopoietic cell differentiation; the physiology is antigen presentation; or the modulating is blocking, and the physiology is lymphoid lineage cell proliferation.

Other embodiments provide methods of modulating a signal to a cell mediated by IL-B50 comprising contacting the cell to an administered agonist or antagonist of IL-B50. These include those wherein the modulating is inhibiting, and the signal is a proliferation signal; the antagonist is a neutralizing antibody to IL-7R α or the R δ 2 subunit or a complex
5 comprising the subunits; the agonist or antagonist is administered in combination with another antagonist or agonist of IL-B50; the agonist or antagonist is administered in combination with a growth factor, cytokine, chemokine, or immune adjuvant; or the contacting is with another anti-proliferative agent or treatment.

Other methods include those of selectively labeling a population of cells, the method
10 comprising contacting the cells with an antibody which binds: IL-7R α ; R δ 2; or a complex comprising one of the subunits; thereby resulting in the identification of cells expressing the subunit or complex. Certain embodiments include those wherein: the contacting results in modulation of STAT activation; the labeling allows purification of IL-7R α or R δ 2 subunit expressing cells; or the labeling allows depletion of IL-7R α or R δ 2 subunit expressing cells.
15 Also provided are populations of cells made by the methods, including those which are prepared by Fluorescent Activated Cell Sorting.

The invention further provides methods of testing a compound for ability to affect receptor-ligand interaction, the method comprising comparing the interaction of a receptor complex comprising IL-7R α and/or R δ 2 subunit with IL-B50 in the presence and absence of
20 the compound. In certain embodiments, the compound is an antibody which binds one of: IL-7R α ; R δ 2 subunit; a receptor comprising IL-7R and/or R δ 2; or IL-B50.

Certain compositions are provided, e.g., an isolated or recombinant protein complex comprising: at least 15 contiguous amino acid residues of SEQ ID NO: 2 and at least 15
contiguous amino acid residues of SEQ ID NO: 4; at least two distinct segments of at least 8
25 contiguous amino acid residues of SEQ ID NO: 2 and at least two distinct segments of at least 8 contiguous amino acid residues of SEQ ID NO: 4; or at least one segment at least 21 contiguous nucleotides of SEQ ID NO: 1 and at least one segment at least 21 contiguous nucleotides of SEQ ID NO: 3. In preferred embodiments, one of the segments of SEQ ID
NO: 2 is from the extracellular portion of the sequence; one of the segments of SEQ ID NO:
30 4 is from the extracellular portion of the sequence; or the polypeptide comprises the mature SEQ ID NO: 2 and the mature SEQ ID NO: 4 sequences.

Nucleic acid embodiments include an isolated or recombinant polynucleotide encoding described components of the complex, wherein: the polynucleotide comprises a deoxyribonucleotide; the polynucleotide comprises a ribonucleotide; or at least one of the segments is operably linked to a promoter.

5 Antibodies are also provided which recognize epitopes presented by the complex, e.g., a binding compound comprising an antigen binding portion from an antibody which binds with selectivity to a polypeptide comprising at least 12 contiguous amino acid residues of SEQ ID NO: 2 and at least 12 contiguous amino acid residues of SEQ ID NO: 4.

10 BRIEF DESCRIPTION OF THE FIGURES

Figures 1A-1E show the nucleotide and amino acid sequences (see SEQ ID NO: 1 and 2) of a primate, e.g., human, IL-7R α ; predicted signal cleavage site indicated.

Figures 2A-2E show the nucleotide and amino acid sequences (see SEQ ID NO: 3 and 4) of a primate, e.g., human, R δ 2; predicted signal cleavage site indicated.

15 Figures 3A and 3B show the amino acid and nucleotide sequences, respectively, of primate, e.g., human, IL-B50; predicted signal cleavage position indicated.

Figures 4A-4E show expression levels of hIL-7R α (Figures 4A and 4B), R δ 2 (hTSLPR, Figures 4C and 4D), and IL-B50 (Figure 4E) in various tissues and cell types. Expression levels were normalized and expressed as femtograms mRNA per 50 ng total
20 cDNA.

Figure 5 shows the induction of TARC by IL-B50. Human CD11c⁺ DC were cultured in the absence or the presence of IL-B50 (50 ng/ml) and the production of TARC was determined in the culture supernatant by ELISA.

25 Figure 6A depicts a culture of sorted CD11c⁺ DC after 24 h in medium alone. DC form small and irregular clumps with a dark center, indicating the presence of dying cells.

Figure 6B depicts a culture of sorted CD11c⁺ DC from the same donor as in Figure 6A, treated with 15 ng/ml of IL-B50. DC form larger and round clumps with fine dendrites visible at the periphery, indicating the maturation of the DC.

30 Figure 7 shows the surface phenotype of CD11c⁺ DC after 24 h of culture with and without (medium alone) IL-B50 and shows the upregulation of HLA-DR, as well as the

costimulatory molecules CD40, CD80 and CD86. Results shown are from one representative of four independent experiments.

Figure 8 shows the surface phenotype of DC after treatment with medium alone, IL-B50, CD40-ligand (CD40L), IL-7 and LPS. IL-B50 is more potent than CD40-ligand and IL-7 in upregulating costimulatory molecules CD40 and CD80.

Figure 9 shows the results of a T cell proliferation assay using CD11c+ DC matured for 24 h in medium or with IL-B50 (15 ng/ml) and cocultured with 5×10^4 allogenic CD4+CD45RA+ naïve T cells at increasing DC/T cell ratios. Proliferation was assessed on day 6 by measuring [3 H]thymidine incorporation. Each point represents the mean [3 H]thymidine incorporation of triplicate cocultures. Vertical bars indicate the SD. DC alone (\square) were used as a control and did not significantly proliferate. Results shown are from one representative of the two independent experiments.

Figure 10 shows the results of a similar experiment as described for Figure 9, using DC matured in medium, IL-B50, CD40-ligand (CD40L), IL-7 and LPS.

Figures 11A-11E show the production of various cytokines (expressed as pg/ml) by naïve CD4 T cells cocultured with DC matured in medium alone, IL-B50, CD40-ligand (CD40L), IL-7 and LPS. Figure 11A shows the effect on the production of IL-4; Figure 11B shows the effect on the production of IL-13; Figure 11C shows the effect on the production of IFN- γ , Figure 11D shows the effect on the production of IL-10 and Figure 11E shows the effect on the production of TNF- α .

Figure 12 shows the effect of DCs treated with medium alone, IL-B50, IL-7, and CD40-ligand on CD8 T cell expansion.

Figure 13 compares expression of perforin by human naïve CD8 T cells induced by DCs treated with medium alone, IL-B50 or CD40-ligand.

Figures 14A-14C show the results of a comparison of IL-B50 with GM-CSF, IL-7, CD40-ligand (CD40L) and medium alone as a control, to stimulate human DCs to produce mRNA for various cytokines and chemokines. Figure 14A shows effects on IL-1 α , IL-1 β , IL-6, IL-12p40 and TNF- α . Figure 14B shows effects on TARC, MDC and MIP3- β . Figure 14C shows effects on MCP-1, MCP-4, Rantes and MIG.

Figure 15 shows the effect of IL-B50 on the induction of IL12p75 protein.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

OUTLINE

- I. General
- 5 II. Activities
- III. Nucleic acids
 - A. encoding fragments, sequence, probes
 - B. mutations, chimeras, fusions
 - C. making nucleic acids
 - 10 D. vectors, cells comprising
- IV. Proteins, Peptides
 - A. fragments, sequence, immunogens, antigens
 - B. muteins
 - C. agonists/antagonists, functional equivalents
 - 15 D. making proteins
- V. Making nucleic acids, proteins
- VI. Antibodies
 - A. polyclonals
 - B. monoclonal, Kd
 - 20 C. anti-idiotypic antibodies
 - D. hybridoma cell lines
- VII. Kits and Methods to quantify ligand/receptor
 - A. ELISA
 - B. assay mRNA encoding
 - 25 C. qualitative/quantitative
 - D. kits
- VIII. Therapeutic compositions, methods
 - A. combination compositions
 - B. unit dose
 - 30 C. administration
- IX. Receptors

I. General

Before the present compositions, formulations, and methods are described, it is to be
 35 understood that this invention is not limited to the particular methods, compositions, and
 cell lines described herein, as such methods, compositions, and cell lines may, of course,
 vary. It is also to be understood that the terminology used herein is for the purpose of
 describing particular embodiments.

As used herein, including the appended claims, singular forms of words such as "a,"
 40 "an," and "the" include their corresponding plural referents unless the context clearly
 dictates otherwise. Thus, e.g., reference to "an organism" includes one or more different
 organisms, reference to "a cell" includes one or more of such cells, and reference to "a

method" includes reference to equivalent steps and methods known to a person of ordinary skill in the art, and so forth.

Unless otherwise defined, technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described
5 herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. Publications, patent applications, patents, and other references discussed above are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the invention
10 is not entitled to antedate any such disclosure by virtue of its prior invention. Publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety including all figures and drawings.

The present invention is based on the discovery of the receptor subunits for the IL-B50 cytokine. This allows advantageous coproduction of the subunits on vectors,
15 production of fusion proteins, and antibody preparations which recognize epitopes resulting from the interaction of the subunit components into a functional unit.

IL-7 biology is reasonably well described. See, e.g., Stoddart, et al. (2000) Immunol. Rev. 175:47-58; Puel and Leonard (2000) Curr. Opin. Immunol. 12:468-473; Akashi, et al. (2000) Curr. Opin. Immunol. 12:144-150; Watanabe, et al. (1999) Immunol. Res. 20:251-259; Waldmann (2000) Ann. Oncol. 11 Suppl 1:101-106; Beverley and
20 Grubeck-Loebenstien (2000) Vaccine 18:1721-1724; Aspinall and Andrew (2000) Vaccine 18:1629-1637; Appasamy (1999) Cytokines Cell Mol. Ther. 5:25-39; Hofmeister, et al. (1999) Cytokine Growth Factor Rev. 10:41-60; Or, et al. (1998) Cytokines Cell Mol. Ther. 4:287-294; Akashi, et al. (1998) Immunol. Rev. 165:13-28; and Offner and Plum (1998)
25 Leuk. Lymphoma 30:87-99. Moreover, since the IL-7 receptor and the IL-B50 receptor share one subunit, the signaling pathways and biology should significantly overlap. This is similar to the GM/IL-3/IL-5 family, which is one of the first groups whose overlapping biologies were explained by the sharing of receptor subunits.

Additionally, recognition of the receptor subunits provides the opportunity to
30 determine cell types and developmental stages where the functional receptor components are coordinately expressed. This provides the opportunity to determine what cell types are likely to respond to ligand, and the resulting biological functions mediated by those cells

provides suggestions as to the physiological effects mediated by the ligand. This leads to better understanding of therapeutic uses of the ligand or blocking ligand:receptor interaction and signaling.

Some of the standard methods applicable are described or referenced, e.g., in
5 Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York; each of which is
10 incorporated herein by reference. See also USSN 09/130,972, which is incorporated herein by reference.

A complete nucleotide (SEQ ID NO: 1) and corresponding amino acid sequence (SEQ ID NO: 2) of a primate, e.g., human, IL-7R α coding segment is shown in Figure 1; similarly for R82 in Figure 2. Figure 3 provides sequence of primate, e.g., human, IL-B50.
15 See USSN 09/399,492, which is incorporated herein by reference.

For the IL-7R α subunit, notable predicted features include, e.g., CK2 phosphorylation sites at about S24-E27, T47-E50, S187-D190, S214-E217, S255-D258, S282-D285, S355-D358, and S423-E426; peroxisomal motif at about S133-L135; PKC phosphorylation sites at about T76-K78, T97-K99, S165-K167, S315-K317, S364-R366, and
20 S373-K375; receptor cyt2 motif at about G195-S201; cAMP kinase motif at about f342-S346; cAMP kinase motif at about R343-S346 or R343-L367; GSK3 motifs at about T129-S133, S201-Y205, T208-n212, S324-p330, T337-f341, S363-S367, and T413-S417; N glycosylation sites at about N29-S31, N45-T47, N131-S133, N162-S164, N212-S214, N213-S215, N275-S277, N353-S355, and N392-T394; tyrosine kinase motif at about E14-Y18;
25 cAMP PK sites at about K94-T97, K84-S87, R343-S346, Cas phos sites at about T208-K210, S215-E217, T262-E264, S331-D333, T337-E339, and S367-D369; cyt C me sites at about T76-F79, C98-I101, Q136-Y139, W244-R247, H259-T261, and C267-P270; histone methylation sites at about F85-L88, Q172-L175, and P270-N273; myristoly sites at about G322-S326, G352-A256, and G389-S392; Phos2 sites at about K184/S187, R343/S345,
30 R343/S346, and R371/S373; and PKC phos sites at about T76-K78, T97-K99, S165-K167, S315-K317, S364-R366, and S373-K375.

Regions of particular interest from the R δ 2 sequence include predictions of, e.g., CK2 phosphorylation sites at about T115-D118, S120-D123, S132-D135, S140-E143, S242-D245, S284-E287, and T297-E300; peroxisomal localization motifs at about S218-F220 and A272-L274; PKC phosphorylation sites at about S76-R78, S99-K101, and S300-R302; tyrosine phosphorylation sites at about R39/D43/Y46 and K166/E169/Y172; cAMP kinase sites at about I77-T81 and R78-T82; Ca⁺⁺ kinase site at about R78-h82; GSK3 sites at about T23-S27, S99-v103, T113-S117, S132-t136, T136-S140, T199-p203, T201-p205, T297-S301, and S301-l305; SigPase sites at about A195-A197 and A278-A280; Tyr kinase site at about D70-Y74; cAMP PK sites at about R78-T81, K96-S99, K157-S160, and K312-S315; Ca⁺⁺ phosphatase sites at about T149-E151 and T309-E311; cyt C Me site at about V235-F238; myristoyl sites at about F3-G7 and G329-T333; N glycosylation sites at about N25-S27, N33-T35, N79-T81, and N147-T149; phos2 sites at about K28/S30, K96/S98, K96/S99, R104/S106, K206/S208, and K312/S315; PKC phosphorylation sites at about S76-R78, S99-K101, and S301-R303; SPKK sites at about S99-h103 and S301-m303; and Tyr Kinase sites at about R39-Y46.

In the IL-B50 sequence, the region from K97-K103 is known to be subject to proteolysis, and mutations may be targeted to that region to protect the ligand from proteolytic degradation. Thus, pharmacokinetic properties of the ligand may be modified, especially for the indications described herein.

Segments with boundaries adjacent these positions will be particularly useful, as will polynucleotides encoding such segments. Mutagenesis in these regions will be used to determine structure-activity relationship, particularly with the receptor, as provided herein.

As used herein, the term "primate IL-7R α " shall be used to describe a protein comprising a protein or peptide segment having or sharing the amino acid sequence shown in Figures 1A-1E (SEQ ID NO:2), or a substantial fragment thereof; but distinct from rodent sequences. The invention also includes protein variations of the IL-7R α allele whose sequence is provided, e.g., a mutein agonist or antagonist. Typically, such agonists or antagonists will exhibit less than about 10% sequence differences, and thus will often have between 1- and 11-fold substitutions, e.g., 2-, 3-, 5-, 7-fold, and others. It also encompasses allelic and other variants, e.g., natural polymorphic variants, of the protein described.

"Natural" as used herein means unmodified by artifice, found, e.g., in natural sources.

Typically, it will bind, when in a functional receptor complex, to ligand with high affinity, e.g., at least about 100 nM, usually better than about 30 nM, preferably better than about 10 nM, and more preferably at better than about 3 nM. The term shall also be used herein to refer to related naturally occurring forms, e.g., alleles, polymorphic variants, and metabolic variants of the primate protein. Corresponding meanings apply to uses of terms related to the R82 sequences.

This invention also encompasses proteins or peptides having amino acid sequence homology with combinations of amino acid sequences presented in Figures 1 and 2. In particular, it will include fusion constructs or proteins comprising segments from both of the sequences provided.

A substantial polypeptide "fragment", or "segment", is a stretch of amino acid residues of at least about 8 amino acids, generally at least 10 amino acids, more generally at least 12 amino acids, often at least 14 amino acids, more often at least 16 amino acids, typically at least 18 amino acids, more typically at least 20 amino acids, usually at least 22 amino acids, more usually at least 24 amino acids, preferably at least 26 amino acids, more preferably at least 28 amino acids, and, in particularly preferred embodiments, at least about 30 or more amino acids. Sequences of segments of different proteins can be compared to one another over appropriate length stretches.

Amino acid sequence homology, or sequence identity, is determined by optimizing residue matches, if necessary, by introducing gaps as required. See, e.g., Needleham, et al., (1970) J. Mol. Biol. 48:443-453; Sankoff, et al., (1983) chapter one in Time Warps, String Edits, and Macromolecules: The Theory and Practice of Sequence Comparison, Addison-Wesley, Reading, MA; and software packages from IntelliGenetics, Mountain View, CA; and the University of Wisconsin Genetics Computer Group (GCG), Madison, WI; each of which is incorporated herein by reference. This changes when considering conservative substitutions as matches. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid; asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. Homologous amino acid sequences are intended to include natural allelic and interspecies variations in the cytokine sequence. Typical homologous proteins or peptides will have from 50-100% homology (if gaps can be introduced), to 60-100% homology (if conservative substitutions are included) with an amino acid sequence segment of Figures 1

or 2. Homology measures will be at least about 70%, generally at least 76%, more generally at least 81%, often at least 85%, more often at least 88%, typically at least 90%, more typically at least 92%, usually at least 94%, more usually at least 95%, preferably at least 96%, and more preferably at least 97%, and in particularly preferred embodiments, at least 98% or more. The degree of homology will vary with the length of the compared segments. Homologous proteins or peptides, such as the allelic variants, will share most biological activities with the embodiments shown in Figures 1 and 2. As used herein, the term "biological activity" is used to describe, without limitation, effects on immune or other cell proliferation, differentiation, induction of cytokines or chemokines (e.g., TARC, PARC, MDC, MIP3-b, etc.), effects on STAT3 and/or STAT5 mediated signal transduction, antigen presentation effects, changes in cell surface molecule expression, and Th2 specific activities.

The terms ligand, agonist, antagonist, and analog of these receptors, particularly the functional complex, include molecules that modulate the characteristic cellular responses to the IL-B50, as well as molecules possessing the more standard structural binding competition features of ligand-receptor interactions, e.g., where the receptor is a natural receptor or an antibody. The cellular responses likely are mediated through binding of IL-B50 to cellular receptors, as described. Also, a ligand is a molecule which serves either as a natural ligand to which said receptor, or an analog thereof, binds, or a molecule which is a functional analog of the natural ligand. The functional analog may be a ligand with structural modifications, or may be a wholly unrelated molecule which has a molecular shape which interacts with the appropriate ligand binding determinants. The ligands may serve as agonists or antagonists, see, e.g., Goodman, et al. (eds.) (1990) Goodman & Gilman's: The Pharmacological Bases of Therapeutics, Pergamon Press, New York.

Rational drug design may also be based upon structural studies of the molecular shapes of a receptor or antibody and other effectors or ligands. Effectors may be other proteins which mediate other functions in response to ligand binding, or other proteins which normally interact with the receptor. One means for determining which sites interact with specific other proteins is a physical structure determination, e.g., x-ray crystallography or 2 dimensional NMR techniques. These will provide guidance as to which amino acid residues form molecular contact regions. For a detailed description of protein structural determination, see, e.g., Blundell and Johnson (1976) Protein Crystallography, Academic Press, New York, which is hereby incorporated herein by reference.

II. Activities

The IL-B50 proteins have a number of different biological activities based on coexpression of IL-7R α and R δ 2, e.g., in the immune system, and include proliferative, developmental, or physiological functions, in particular of lymphoid lineage cells, e.g., macrophages or dendritic cells. The IL-B50 proteins are homologous to other IL-7 ligand family proteins, but each have structural differences. For example, human IL-B50 shows 43% amino acid sequence identity to mouse TSLP. Additionally, the human receptor subunit R δ 2 displays 39% amino acid sequence identity to mouse TSLPR.

The mouse IL-B50 molecule has the ability to stimulate TARC production and various Th2 specific cytokines. The signaling pathway seems to use STAT3 and/or STAT5, and sends proliferation or differentiation signals. Differentiation tends to result in limitation of proliferation, and vice versa. Differentiation typically results in changes in cell surface marker expression.

As shown herein, human IL-B50 improves dendritic cell survival in cultures, upregulates the expression of costimulatory molecules and adhesion molecules, including HLA-DR, CD40, CD80, CD86, CD11a, CD18 and CD83, induces dendritic cells to produce the chemokines TARC, PARC and MDC, and strongly promotes the capacity of dendritic cells to induce naïve T cells to proliferate and to produce cytokines IL-4, IL-13, and TNF- α . Additionally, IL-B50 has a synergistic effect with CD40-ligand and LPS in activated dendritic cells to upregulate costimulatory molecules CD40, CD80 and CD86. IL-B50 also strongly enhances CD40-ligand-induced production of IL-12 by dendritic cells.

TARC, PARC and MDC are all notably ligands for CCR4, a chemokine receptor predominantly found on Th2-type lymphocytes. Thus, IL-B50 can activate myeloid cells, such as monocytes, to release chemokines that may attract effector cells with a Th2 phenotype. As shown in the examples below, IL-B50-induced expression of TARC was very strong in the CD11c⁺ subset of DCs. This subset, representing less than 1% of mononuclear cells in the blood, normally differentiates into mature DCs in response to inflammatory stimuli. The expression of TARC in these cells was accompanied by a dramatic enhancement of their maturation as evidenced by the strong induction of the costimulatory molecules CD40 and CD80. These results indicate that this DC subset

stimulated with IL-B50 could be a potent inducer of primary T-cell-mediated immune responses. Indeed, CD11+ DCs cultured in the presence of IL-B50 are much more potent in their capacity to elicit the proliferation of naïve T cells as compared to DC cultured in medium.

5 Dendritic cells are professional antigen presenting cells, which are capable of inducing primary antigen-specific T cell-mediated immune responses. Dendritic cells play a critical role in initiating immune responses against tumors and infectious microorganisms. Dendritic cells are also involved in autoimmune diseases, allergic diseases, graft-versus-host disease and rejection of solid organ transplants. Therefore, enhancing dendritic cell
10 function allows for treatment of tumors and infectious diseases. Similarly, blocking dendritic cell function provides therapies for autoimmune diseases, allergic diseases, graft-versus-host diseases and transplantation associated rejection.

Thus, IL-B50 may be used in enhancing dendritic cell function in treating cancers and infectious diseases and IL-B50 antagonists may be used in blocking the function of
15 dendritic cells in treating autoimmune diseases, allergic diseases, graft-versus-host diseases and transplantation associated rejection. The elucidation of the IL-B50 receptor subunits, therefore, allows for the identification of agonists and antagonists of ILB-50 for use in treating the aforementioned diseases.

The present disclosure also describes new assays for activities described for these
20 molecules. Corresponding activities should be found in other mammalian systems, including primates. The new IL-7-like molecules produced by recombinant means exhibit a biological activity of modulating lymphoid lineage cells. Furthermore, there is substantial likelihood of synergy with other IL-7 related agonists or antagonists. It is likely that the receptors, which are expected to include multiple different polypeptide chains, exhibit
25 species specificity for their corresponding ligands.

III. Nucleic Acids

This invention contemplates use of isolated nucleic acid or fragments, e.g., which encode these or closely related proteins, or fragments thereof, e.g., fusion proteins or
30 coordinately expressed or combination expression constructs.

The term "isolated nucleic acid or fragments" as used herein means a nucleic acid, e.g., a DNA or RNA molecule, that is not immediately contiguous with sequences present in

the naturally occurring genome of the organism from which it is derived. Thus, the term describes, e.g., a nucleic acid that is incorporated into a vector, such as a plasmid or viral vector; a nucleic acid that is incorporated into the genome of a heterologous cell (or the genome of homologous cell, but at a site different from that at which it normally occurs);
5 and a nucleic acid that exists as a separate molecule, e.g., a DNA fragment produced by PCR amplification or restriction enzyme digestion, or an RNA molecule produced by in vitro transcription. The term also describes a recombinant (e.g., genetically engineered) nucleic acid that forms part of a hybrid gene encoding additional polypeptide sequences that can be used, e.g., in the production of a fusion protein. In addition, this invention embodies
10 any engineered or nucleic acid molecule created by artifice that encodes a biologically active protein or polypeptide having characteristic IL-B50 receptor activity. Typically, the nucleic acid is capable of hybridizing, under appropriate conditions, with nucleic acid sequence segments shown in Figures 1 and 2. Further, this invention covers the use of isolated or recombinant nucleic acid, or fragments thereof, which encode proteins having
15 fragments which are homologous to the newly disclosed receptor complex proteins. The isolated nucleic acids can have the respective regulatory sequences in the 5' and 3' flanks, e.g., promoters, enhancers, poly-A addition signals, and others from the natural gene.

An "isolated" nucleic acid is a nucleic acid, e.g., an RNA, DNA, or a mixed polymer, which is substantially pure, e.g., separated from other components which naturally
20 accompany a native sequence, such as ribosomes, polymerases, and flanking genomic sequences from the originating species. The term embraces a nucleic acid sequence which has been removed from its naturally occurring environment, and includes recombinant or cloned DNA isolates, which are thereby distinguishable from naturally occurring compositions, and chemically synthesized analogs or analogs biologically synthesized by
25 heterologous systems. A substantially pure molecule includes isolated forms of the molecule, either completely or substantially pure.

An isolated nucleic acid will generally be a homogeneous composition of molecules, but will, in some embodiments, contain heterogeneity, preferably minor. This heterogeneity is typically found at the polymer ends or portions not critical to a desired biological function
30 or activity.

A "recombinant" nucleic acid is defined either by its method of production or its structure. In reference to its method of production, e.g., a product made by a process, the

process is use of recombinant nucleic acid techniques, e.g., involving human intervention in the nucleotide sequence. Typically this intervention involves in vitro manipulation, although under certain circumstances it may involve more classical animal breeding techniques. Alternatively, it can be a nucleic acid made by generating a sequence
5 comprising fusion of two fragments which are not naturally contiguous to each other, but is meant to exclude products of nature, e.g., naturally occurring mutants as found in their natural state. Thus, e.g., products made by transforming cells with any unnaturally occurring vector is encompassed, as are nucleic acids comprising sequence derived using any synthetic oligonucleotide process. Such a process is often done to replace a codon with
10 a redundant codon encoding the same or a conservative amino acid, while typically introducing or removing a restriction enzyme sequence recognition site. Alternatively, the process is performed to join together nucleic acid segments of desired functions to generate a single genetic entity comprising a desired combination of functions not found in the commonly available natural forms, e.g., encoding a fusion protein. Restriction enzyme
15 recognition sites are often the target of such artificial manipulations, but other site specific targets, e.g., promoters, DNA replication sites, regulation sequences, control sequences, or other useful features may be incorporated by design. A similar concept is intended for a recombinant, e.g., fusion, polypeptide. This will include a dimeric repeat. Specifically included are synthetic nucleic acids which, by genetic code redundancy, encode similar
20 combination polypeptides to fragments of the receptor subunits and fusions of sequences from various different receptors or related molecules, e.g., growth factors.

A "fragment" in a nucleic acid context is a contiguous segment of at least about 17 nucleotides, generally at least 21 nucleotides, more generally at least 25 nucleotides, ordinarily at least 30 nucleotides, more ordinarily at least 35 nucleotides, often at least 39
25 nucleotides, more often at least 45 nucleotides, typically at least 50 nucleotides, more typically at least 55 nucleotides, usually at least 60 nucleotides, more usually at least 66 nucleotides, preferably at least 72 nucleotides, more preferably at least 79 nucleotides, and in particularly preferred embodiments will be at least 85 or more nucleotides including, e.g., 100, 150, 200, 250, etc. Typically, fragments of different genetic sequences can be
30 compared to one another over appropriate length stretches, particularly defined segments such as the domains described below.

A nucleic acid which codes for an IL-B50 receptor complex will be particularly useful to identify genes, mRNA, and cDNA species which code for itself or closely related proteins, as well as DNAs which code for polymorphic, allelic, or other genetic variants, e.g., from different individuals or related species. Preferred probes for such screens are those regions of the interleukin which are conserved between different polymorphic variants or which contain nucleotides which lack specificity, and will preferably be full length or nearly so. In other situations, polymorphic variant specific sequences will be more useful.

This invention further covers recombinant nucleic acid molecules and fragments having a nucleic acid sequence identical to or highly homologous to the isolated DNA set forth herein. In particular, the sequences will often be operably linked to DNA segments which control transcription, translation, and DNA replication. These additional segments typically assist in expression of the desired nucleic acid segment.

Homologous nucleic acid sequences, when compared to one another or to the sequences shown in Figures 1 or 2, exhibit significant similarity. The standards for homology in nucleic acids are either measures for homology generally used in the art by sequence comparison or based upon hybridization conditions. Comparative hybridization conditions are described in greater detail below.

Substantial identity in the nucleic acid sequence comparison context means either that the segments, or their complementary strands, when compared, are identical when optimally aligned, with appropriate nucleotide insertions or deletions, in at least about 60% of the nucleotides, generally at least 66%, ordinarily at least 71%, often at least 76%, more often at least 80%, usually at least 84%, more usually at least 88%, typically at least 91%, more typically at least about 93%, preferably at least about 95%, more preferably at least about 96 to 98% or more, and in particular embodiments, as high at about 99% or more of the nucleotides, including, e.g., segments encoding structural domains such as the segments described below. Alternatively, substantial identity will exist when the segments will hybridize under selective hybridization conditions, to a strand or its complement, typically using a sequence derived from the sequences depicted in Figures 1 and 2. Typically, selective hybridization will occur when there is at least about 55% homology over a stretch of at least about 14 nucleotides, more typically at least about 65%, preferably at least about 75%, and more preferably at least about 90%. See, Kanehisa (1984) Nuc. Acids Res. 12:203-213. The length of homology comparison, as described, may be over longer

stretches, and in certain embodiments will be over a stretch of at least about 17 nucleotides, generally at least about 20 nucleotides, ordinarily at least about 24 nucleotides, usually at least about 28 nucleotides, typically at least about 32 nucleotides, more typically at least about 40 nucleotides, preferably at least about 50 nucleotides, and more preferably at least
5 about 75 to 100 or more nucleotides.

Stringent conditions, in referring to homology in the hybridization context, will be stringent combined conditions of salt, temperature, organic solvents, and other parameters typically controlled in hybridization reactions. Stringent temperature conditions will usually include temperatures in excess of about 30 degrees C, more usually in excess of about 37
10 degrees C, typically in excess of about 45 degrees C, more typically in excess of about 55 degrees C, preferably in excess of about 65 degrees C, and more preferably in excess of about 70 degrees C. Stringent salt conditions will ordinarily be less than about 500 mM, usually less than about 400 mM, more usually less than about 300 mM, typically less than about 200 mM, preferably less than about 100 mM, and more preferably less than about 80
15 mM, even down to less than about 20 mM. Certain detergents or destabilizing reagents may be added, e.g., formamide at 50%, etc. However, the combination of parameters is much more important than the measure of any single parameter. See, e.g., Wetmur and Davidson (1968) J. Mol. Biol. 31:349-370, which is hereby incorporated herein by reference.

The isolated DNA can be readily modified by nucleotide substitutions, nucleotide
20 deletions, nucleotide insertions, and inversions of nucleotide stretches. These modifications result in novel DNA sequences which encode this protein or its derivatives. These modified sequences can be used to produce mutant proteins (muteins) or to enhance the expression of variant species. Enhanced expression may involve gene amplification, increased transcription, increased translation, and other mechanisms. Such mutant receptor-like
25 derivatives include predetermined or site-specific mutations of the protein or its fragments, including silent mutations using genetic code degeneracy. "Mutant IL-B50 receptor" as used herein encompasses a polypeptide otherwise falling within the homology definition of the receptor as set forth above, but having an amino acid sequence which differs from that of other IL-7 receptor-like proteins as found in nature, whether by way of deletion,
30 substitution, or insertion. In particular, "site specific mutant IL-B50 receptor" encompasses a protein having substantial homology with a protein shown in Figures 1 and 2, and typically shares most of the biological activities of the form disclosed herein.

Although site specific mutation sites are predetermined, mutants need not be site specific. Mammalian IL-B50 receptor mutagenesis can be achieved by making amino acid insertions or deletions in the gene, coupled with expression. Substitutions, deletions, insertions, or any combinations may be generated to arrive at a final construct. Insertions
5 include amino- or carboxy- terminal fusions. Random mutagenesis can be conducted at a target codon and the expressed mammalian IL-B50 receptor mutants can then be screened for the desired activity. Methods for making substitution mutations at predetermined sites in DNA having a known sequence are well known in the art, e.g., by M13 primer mutagenesis. See also Sambrook, et al. (1989) and Ausubel, et al. (1987 and periodic
10 Supplements).

The mutations in the DNA normally should not place coding sequences out of reading frames and preferably will not create complementary regions that could hybridize to produce secondary mRNA structure such as loops or hairpins.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra-
15 Letts, 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Polymerase chain reaction (PCR) techniques can often be applied in mutagenesis.
20 Alternatively, mutagenesis primers are commonly used methods for generating defined mutations at predetermined sites. See, e.g., Innis, et al. (eds. 1990) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (eds. 1995) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY.

25

IV. Proteins, Peptides

As described above, the present invention encompasses primate IL-B50 receptor, e.g., whose sequences are disclosed in Figures 1 and 2, and described above. Allelic and other variants are also contemplated, including, e.g., fusion proteins combining portions of
30 such sequences with others, including epitope tags and functional domains.

The present invention also provides recombinant proteins, e.g., heterologous fusion proteins using segments from these primate proteins. A heterologous fusion protein is a

fusion of proteins or segments which are naturally not normally fused in the same manner. Thus, the fusion product of a growth factor receptor with a cytokine receptor is a continuous protein molecule having sequences fused in a typical peptide linkage, typically made as a single translation product and exhibiting properties, e.g., antigenicity, derived from each source peptide. A similar concept applies to heterologous nucleic acid sequences.

In addition, new constructs may be made from combining similar functional or structural domains from other related proteins, e.g., growth factors or other cytokines. For example, receptor-binding or other segments may be "swapped" between different new fusion polypeptides or fragments. See, e.g., Cunningham, et al. (1989) Science 243:1330-1336; and O'Dowd, et al. (1988) J. Biol. Chem. 263:15985-15992, each of which is incorporated herein by reference. Thus, new chimeric polypeptides exhibiting new combinations of specificities will result from the functional linkage of receptor-binding specificities. For example, the ligand binding domains from other related receptor molecules may be added or substituted for other domains of these or related proteins. The resulting protein will often have hybrid function and properties. For example, a fusion protein may include a labeling epitope which may serve to provide diagnostic locatability of the fusion protein for histology or other methods.

Candidate fusion partners and sequences can be selected from various sequence data bases, e.g., GenBank, c/o IntelliGenetics, Mountain View, CA; and BCG, University of Wisconsin Biotechnology Computing Group, Madison, WI, which are each incorporated herein by reference.

"Derivatives" of the mammalian IL-B50 receptor include amino acid sequence mutants, glycosylation variants, metabolic derivatives and covalent or aggregative conjugates with other chemical moieties. Covalent derivatives can be prepared by linkage of functionalities to groups which are found in the IL-B50 receptor amino acid side chains or at the N- or C- termini, e.g., by means which are well known in the art. These derivatives can include, without limitation, aliphatic esters or amides of the carboxyl terminus, or of residues containing carboxyl side chains, O-acyl derivatives of hydroxyl group-containing residues, and N-acyl derivatives of the amino terminal amino acid or amino-group containing residues, e.g., lysine or arginine. Acyl groups are selected from the group of alkyl-moieties including C3 to C18 normal alkyl, thereby forming alkanoyl aroyl species.

In particular, glycosylation alterations are included, e.g., made by modifying the glycosylation patterns of a polypeptide during its synthesis and processing, or in further processing steps. Particularly preferred means for accomplishing this are by exposing the polypeptide to glycosylating enzymes derived from cells which normally provide such processing, e.g., mammalian glycosylation enzymes. Deglycosylation enzymes are also contemplated. Also embraced are versions of the same primary amino acid sequence which have other minor modifications, including phosphorylated amino acid residues, e.g., phosphotyrosine, phosphoserine, or phosphothreonine.

A major group of derivatives are covalent conjugates of the interleukin or fragments thereof with other proteins of polypeptides. These derivatives can be synthesized in recombinant culture such as N- or C-terminal fusions or by the use of agents known in the art for their usefulness in cross-linking proteins through reactive side groups. Preferred derivatization sites with cross-linking agents are at free amino groups, carbohydrate moieties, and cysteine residues.

Fusion polypeptides between the receptors and other homologous or heterologous proteins are also provided. Homologous polypeptides may be fusions between different receptors, resulting in, for instance, a hybrid protein exhibiting binding specificity for multiple different ligands, or a receptor which may have broadened or weakened specificity of binding to its ligand. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. Typical examples are fusions of a reporter polypeptide, e.g., luciferase, with a segment or domain of a receptor, e.g., a ligand-binding segment, so that the presence or location of a desired receptor may be easily determined. See, e.g., Dull, et al., U.S. Patent No. 4,859,609, which is hereby incorporated herein by reference. Other gene fusion partners include glutathione-S-transferase (GST), bacterial β -galactosidase, trpE, Protein A, β -lactamase, alpha amylase, alcohol dehydrogenase, yeast alpha mating factor, or other cytokine receptors. See, e.g., Godowski, et al. (1988) Science 241:812-816.

The phosphoramidite method described by Beaucage and Carruthers (1981) Tetra. Letts. 22:1859-1862, will produce suitable synthetic DNA fragments. A double stranded fragment will often be obtained either by synthesizing the complementary strand and annealing the strand together under appropriate conditions or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Such polypeptides may also have amino acid residues which have been chemically modified by phosphorylation, sulfonation, biotinylation, PEGylation, or the addition or removal of other moieties, particularly those which have molecular shapes similar to phosphate groups. In some embodiments, the modifications will be useful labeling reagents, or serve as purification targets, e.g., affinity ligands.

Fusion proteins will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, e.g., in Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual (2d ed.), Vols. 1-3, Cold Spring Harbor Laboratory, and Ausubel, et al. (eds. 1987 and periodic supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York, which are each incorporated herein by reference. Techniques for synthesis of polypeptides are described, e.g., in Merrifield (1963) J. Amer. Chem. Soc. 85:2149-2156; Merrifield (1986) Science 232: 341-347; and Atherton, et al. (1989) Solid Phase Peptide Synthesis: A Practical Approach, IRL Press, Oxford; each of which is incorporated herein by reference. See also Dawson, et al. (1994) Science 266:776-779 for methods to make larger polypeptides.

The term "substantially purified" as used herein refers to a molecule, such as a peptide that is substantially free of other proteins, lipids, carbohydrates, nucleic acids, or other biological materials with which it is naturally associated. For example, a substantially pure molecule, such as a polypeptide, can be at least 60%, by dry weight, the molecule of interest. One skilled in the art can purify complexes comprising the described sequences using standard protein purification methods and the purity of the polypeptides can be determined using standard methods including, e.g., polyacrylamide gel electrophoresis (e.g., SDS-PAGE), column chromatography (e.g., high performance liquid chromatography (HPLC)), and amino-terminal amino acid sequence analysis.

Changes in the amino acid sequence of the receptor components of the complex are contemplated in the present invention. The IL-7R α or R δ 2 can be altered by changing the nucleic acid sequence encoding the proteins. Preferably, only conservative amino acid alterations are undertaken, using amino acids that have the same or similar properties. Illustrative amino acid substitutions include the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine

or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine, glutamine, or glutamate; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; valine to isoleucine or leucine.

5 Additionally, other variants and fragments of receptor complexes can be used in the present invention. Variants include analogs, homologues, derivatives, muteins, and mimetics of the receptor complexes or fragments that retain or enhance the ability to block binding between IL-B50 and a target receptor. The variants and fragments can be generated directly from sequences provided, by chemical modification, by proteolytic enzyme
10 digestion, or by combinations thereof. Additionally, genetic engineering techniques, as well as methods of synthesizing polypeptides directly from amino acid residues, can be employed.

Non-peptide compounds that mimic the binding and function of IL-B50 ("mimetics") can be produced by the approach outlined in Saragovi, et al. (1991) Science
15 253:792-95. Mimetics are molecules which mimic elements of protein secondary structure. See, e.g., Johnson et al., "Peptide Turn Mimetics," in Pezzuto, et al. (eds. 1993) Biotechnology and Pharmacy, Chapman and Hall, New York. The underlying rationale behind the use of peptide mimetics is that the peptide backbone of proteins exists chiefly to orient amino acid side chains in such a way as to facilitate molecular interactions. For the
20 purposes of the present invention, appropriate mimetics can be considered to be the equivalent of IL-B50 itself.

Variants and fragments also can be created by recombinant techniques employing genomic or cDNA cloning methods. Site-specific and region-directed mutagenesis techniques can be employed. See, e.g., vol. 1, ch. 8 in Ausubel, et al. (eds. 1989 and
25 periodic updates) Current Protocols in Molecular Biology Wiley and Sons; and Oxender and Fox (eds.) Protein Engineering Liss, Inc. In addition, linker-scanning and PCR-mediated techniques can be employed for mutagenesis. See, e.g., Erlich (ed. 1989) PCR Technology Stockton Press. Protein sequencing, structure and modeling approaches for use with any of the above techniques are disclosed, e.g., in Oxender and Fox (eds.) Protein
30 Engineering Liss, Inc; and Ausubel, et al. (eds. 1989 and periodic updates) Current Protocols in Molecular Biology Wiley and Sons.

This invention also contemplates the use of derivatives of IL-B50 receptor other than variations in amino acid sequence or glycosylation. Such derivatives may involve covalent or aggregative association with chemical moieties. These derivatives generally fall into three classes: (1) salts, (2) side chain and terminal residue covalent modifications, and (3) adsorption complexes, e.g., with cell membranes. Such covalent or aggregative derivatives are useful as immunogens, as reagents in immunoassays, in enhancing pharmacokinetic properties of the protein, or in purification methods such as for affinity purification of a receptor or other binding molecule, e.g., an antibody. For example, an IL-B50 receptor can be immobilized by covalent bonding to a solid support such as cyanogen bromide-activated SEPHAROSE, by methods which are well known in the art, or adsorbed onto polyolefin surfaces, with or without glutaraldehyde cross-linking, for use in the assay or purification of IL-B50 receptor, antibodies, or other similar molecules. The IL-B50 can also be labeled with a detectable group, e.g., radio-iodinated by the chloramine T procedure, covalently bound to rare earth chelates, or conjugated to another fluorescent moiety for use in diagnostic assays.

A receptor complex of this invention can be used as an immunogen for the production of antisera or antibodies specific, e.g., for the complex comprising both protein components. The purified interleukin can be used to screen monoclonal antibodies or antigen-binding fragments prepared by immunization with various forms of impure preparations containing the protein. In particular, the term "antibodies" also encompasses antigen binding fragments of natural antibodies. The purified interleukin can also be used as a reagent to detect any antibodies generated in response to the presence of elevated levels of expression, or immunological disorders which lead to antibody production to the endogenous cytokine. Additionally, receptor complex fragments may also serve as immunogens to produce the antibodies of the present invention, as described immediately below. For example, this invention contemplates antibodies having binding affinity to or being raised against a complex comprising the amino acid sequences shown in Figures 1 and 2, fragments thereof, or homologous peptides. In particular, this invention contemplates antibodies having binding affinity to, or having been raised against, specific fragments which are predicted to be, or actually are, exposed at the exterior protein surface of the native receptor.

The blocking of physiological response to the ligand may result from the inhibition of binding of the ligand to the receptor, likely through competitive inhibition. Thus, in vitro assays of the present invention will often use antibodies or receptor binding segments of these antibodies, or fragments attached to solid phase substrates. These assays will also
5 allow for the diagnostic determination of the effects of either binding region mutations and modifications, or receptor mutations and modifications.

This invention also contemplates the use of competitive drug screening assays, e.g., where neutralizing antibodies to the receptor complex or fragments compete with a test compound for binding. In this manner, the neutralizing antibodies or fragments can be used
10 to detect the presence of a polypeptide which shares one or more binding sites to a receptor and can also be used to occupy binding sites on a receptor that might otherwise bind a cytokine, e.g., IL-B50.

V. Making Nucleic Acids and Protein

DNA which encodes the proteins or fragments thereof can be obtained by chemical
15 synthesis, screening cDNA libraries, or by screening genomic libraries prepared from a wide variety of cell lines or tissue samples. Natural sequences can be isolated using standard methods and the sequences provided herein, e.g., in Figures 1 and 2. Other species counterparts, e.g., primate, can be identified by hybridization techniques, or by various PCR
20 techniques, combined with or by searching in sequence databases.

These DNAs can be expressed in a wide variety of host cells for the synthesis and formation of the functional receptor complex, or fragments, which can in turn, e.g., be used to generate polyclonal or monoclonal antibodies; for binding studies; for construction and expression of modified agonist/antagonist molecules; for structure/function studies; and for
25 screening studies. A variant or its fragments can be expressed in host cells that are transformed or transfected with appropriate expression vectors. These molecules can be substantially free of protein or cellular contaminants, other than those derived from the recombinant host, and therefore are particularly useful in pharmaceutical compositions when combined with a pharmaceutically acceptable carrier and/or diluent. The proteins, or
30 portions thereof, may be expressed as fusions with other proteins or using combining motifs, e.g., leucine zippers, to form a soluble receptor complex.

Expression vectors are typically self-replicating DNA or RNA constructs containing the desired receptor gene or its fragments, usually operably linked to suitable genetic control elements that are recognized in a suitable host cell. These control elements are capable of effecting expression within a suitable host. The specific type of control elements necessary to effect expression will depend upon the eventual host cell used. Generally, the genetic control elements can include a prokaryotic promoter system or a eukaryotic promoter expression control system, and typically include a transcriptional promoter, an optional operator to control the onset of transcription, transcription enhancers to elevate the level of mRNA expression, a sequence that encodes a suitable ribosome binding site, and sequences that terminate transcription and translation. Expression vectors also usually contain an origin of replication that allows the vector to replicate independently of the host cell.

The vectors of this invention include those which contain DNA which encodes these proteins, as described, or a fragment comprising sequence from both subunits, typically encoding a functionally, e.g., biologically active, equivalent polypeptide. The DNA can be under the control of a viral promoter and can encode a selection marker. This invention further contemplates use of such expression vectors which are capable of expressing eukaryotic cDNA coding for such a protein or complex in a prokaryotic or eukaryotic host, where the vector is compatible with the host and where the eukaryotic cDNA coding for the receptor is inserted into the vector such that growth of the host containing the vector expresses the cDNA in question. Usually, expression vectors are designed for stable replication in their host cells or for amplification to greatly increase the total number of copies of the desirable gene per cell. It is not always necessary to require that an expression vector replicate in a host cell, e.g., it is possible to effect transient expression of the receptor or its fragments in various hosts using vectors that do not contain a replication origin that is recognized by the host cell. It is also possible to use vectors that cause integration of the protein encoding portion or its fragments into the host DNA by recombination, e.g., downstream of heterologous promoters, or integration of heterologous promoters upstream from endogenous genes.

Vectors, as used herein, comprise plasmids, viruses, bacteriophage, integratable DNA fragments, and other vehicles, e.g., which may enable the integration of DNA fragments into the genome of the host. Expression vectors are specialized vectors which contain genetic control elements that effect expression of operably linked genes. Plasmids

are the most commonly used form of vector but other forms of vectors which serve an equivalent function and which are, or become, known in the art are suitable for use herein. See, e.g., Pouwels, et al. (1985 and Supplements) Cloning Vectors: A Laboratory Manual, Elsevier, N.Y., and Rodriguez, et al. (eds. 1988) Vectors: A Survey of Molecular Cloning Vectors and Their Uses, Butterworth, Boston, which are incorporated herein by reference.

Transformed cells are cells, preferably mammalian, that have been transformed, transfected, or infected with vectors, e.g., constructed using recombinant DNA techniques. Transformed host cells usually express the desired protein or its fragments, but for purposes of cloning, amplifying, and manipulating its DNA, do not need to express the subject protein. This invention further contemplates culturing transformed cells in a nutrient medium, thus permitting the cells to express the protein, including production of soluble receptor complexes or ligand.

For purposes of this invention, nucleic acid sequences are operably linked when they are functionally related to each other. For example, DNA for a pre-sequence or secretory leader is operably linked to a polypeptide if it is expressed as a pre-protein or participates in directing the polypeptide to the cell membrane or in secretion of the polypeptide. A promoter is operably linked to a coding sequence if it promotes the transcription of the polypeptide; a ribosome binding site is operably linked to a coding sequence if it is positioned to induce translation. Usually, operably linked means contiguous and in reading frame, however, certain genetic elements such as repressor genes are not contiguously linked but still bind to operator sequences that in turn control expression.

Suitable host cells include prokaryotes, lower eukaryotes, and higher eukaryotes. Prokaryotes include both gram negative and gram positive organisms, e.g., E. coli and B. subtilis. Lower eukaryotes include yeasts, e.g., S. cerevisiae and Pichia species, and species of the genus Dictyostelium. Higher eukaryotes include established tissue culture cell lines from animal cells, both of non-mammalian origin, e.g., insect cells, and birds, and of mammalian origin, e.g., human, primates, and rodents.

Prokaryotic host-vector systems include a wide variety of vectors for many different species. As used herein, E. coli and its vectors will be used generically to include equivalent vectors used in other prokaryotes. A common vector for amplifying DNA is pBR322 or many of its derivatives. Vectors that can be used to express the receptor or its fragments include, but are not limited to, such vectors as those containing the lac promoter

(pUC-series); trp promoter (pBR322-trp); Ipp promoter (the pIN-series); lambda-pP or pR promoters (pOTS); or hybrid promoters such as ptac (pDR540). See Brosius, et al. (1988) "Expression Vectors Employing Lambda-, trp-, lac-, and Ipp-derived Promoters", in Rodriguez and Denhardt (eds.) Vectors: A Survey of Molecular Cloning Vectors and Their
5 Uses, Butterworth, Boston, Chapter 10, pp. 205-236, which is incorporated herein by reference.

Lower eukaryotes, e.g., yeasts and Dictyostelium, may be transformed with IL-B50 receptor sequence containing vectors. For purposes of this invention, the most common lower eukaryotic host is the baker's yeast, Saccharomyces cerevisiae. It will be used to
10 generically represent lower eukaryotes although a number of other strains and species are also available. Yeast vectors typically consist of a replication origin (unless of the integrating type), a selection gene, a promoter, DNA encoding the receptor or its fragments, and sequences for translation termination, polyadenylation, and transcription termination. Suitable expression vectors for yeast include such constitutive promoters as
15 3-phosphoglycerate kinase and various other glycolytic enzyme gene promoters or such inducible promoters as the alcohol dehydrogenase 2 promoter or metallothionine promoter. Suitable vectors include derivatives of the following types: self-replicating low copy number (such as the YRp-series), self-replicating high copy number (such as the YEp-series); integrating types (such as the YIp-series), or mini-chromosomes (such as the
20 YCp-series).

Higher eukaryotic tissue culture cells are normally the preferred host cells for expression of the functionally active interleukin protein. In principle, any higher eukaryotic tissue culture cell line is workable, e.g., insect baculovirus expression systems, whether from an invertebrate or vertebrate source. However, mammalian cells are preferred.
25 Transformation or transfection and propagation of such cells has become a routine procedure. Examples of useful cell lines include HeLa cells, Chinese hamster ovary (CHO) cell lines, baby rat kidney (BRK) cell lines, insect cell lines, bird cell lines, and monkey (COS) cell lines. Expression vectors for such cell lines usually include an origin of replication, a promoter, a translation initiation site, RNA splice sites (if genomic DNA is
30 used), a polyadenylation site, and a transcription termination site. These vectors also usually contain a selection gene or amplification gene. Suitable expression vectors may be plasmids, viruses, or retroviruses carrying promoters derived, e.g., from such sources as

from adenovirus, SV40, parvoviruses, vaccinia virus, or cytomegalovirus. Representative examples of suitable expression vectors include pCDNA1; pCD, see Okayama, et al. (1985) Mol. Cell Biol. 5:1136-1142; pMC1neo PolyA, see Thomas, et al. (1987) Cell 51:503-512; and a baculovirus vector such as pAC 373 or pAC 610.

5 For secreted proteins, an open reading frame usually encodes a polypeptide that consists of a mature or secreted product covalently linked at its N-terminus to a signal peptide. The signal peptide is typically cleaved prior to secretion of the mature, or active, polypeptide. The cleavage site can be predicted with a high degree of accuracy from empirical rules, e.g., von-Heijne (1986) Nucleic Acids Research 14:4683-4690, and the
10 precise amino acid composition of the signal peptide does not appear to be critical to its function. See, e.g., Randall, et al. (1989) Science 243:1156-1159; and Kaiser et al. (1987) Science 235:312-317.

 It will often be desired to express these polypeptides in a system which provides a specific or defined glycosylation pattern. In this case, the usual pattern will be that provided
15 naturally by the expression system. However, the pattern will be modifiable by exposing the polypeptide, e.g., an unglycosylated form, to appropriate glycosylating proteins introduced into a heterologous expression system. For example, the receptor genes may be co-transformed with one or more genes encoding mammalian or other glycosylating enzymes. Using this approach, certain mammalian glycosylation patterns will be achievable
20 in prokaryote or other cells.

 The source of receptor complex can be a eukaryotic or prokaryotic host expressing recombinant receptor subunit DNA, such as is described above. The source can also be a cell line such as mouse Swiss 3T3 fibroblasts, but other mammalian cell lines are also contemplated by this invention, with the preferred cell line being from the human species.

25 Now that the entire sequence and components of the functional receptor complex is known, the human receptor complex, fragments, or derivatives thereof can be prepared by conventional processes for synthesizing peptides. These include processes such as are described in Stewart and Young (1984) Solid Phase Peptide Synthesis, Pierce Chemical Co., Rockford, IL; Bodanszky and Bodanszky (1984) The Practice of Peptide Synthesis,
30 Springer-Verlag, New York; and Bodanszky (1984) The Principles of Peptide Synthesis, Springer-Verlag, New York; all of each which are incorporated herein by reference. For example, an azide process, an acid chloride process, an acid anhydride process, a mixed

anhydride process, an active ester process (e.g., p-nitrophenyl ester, N-hydroxysuccinimide ester, or cyanomethyl ester), a carbodiimidazole process, an oxidative-reductive process, or a dicyclohexylcarbodiimide (DCCD)/additive process can be used. Solid phase and solution phase syntheses are both applicable to the foregoing processes.

5 The receptor subunit proteins, fragments, or derivatives are suitably prepared in accordance with the above processes as typically employed in peptide synthesis, generally either by a so-called stepwise process which comprises condensing an amino acid to the terminal amino acid, one by one in sequence, or by coupling peptide fragments to the terminal amino acid. Amino groups that are not being used in the coupling reaction
10 typically must be protected to prevent coupling at an incorrect location.

 If a solid phase synthesis is adopted, the C-terminal amino acid is bound to an insoluble carrier or support through its carboxyl group. The insoluble carrier is not particularly limited as long as it has a binding capability to a reactive carboxyl group. Examples of such insoluble carriers include halomethyl resins, such as chloromethyl resin or
15 bromomethyl resin, hydroxymethyl resins, phenol resins, tert-alkyloxycarbonylhydrazidated resins, and the like.

 An amino group-protected amino acid is bound in sequence through condensation of its activated carboxyl group and the reactive amino group of the previously formed peptide or chain, to synthesize the peptide step by step. After synthesizing the complete sequence,
20 the peptide is split off from the insoluble carrier to produce the peptide. This solid-phase approach is generally described by Merrifield, et al. (1963) in J. Am. Chem. Soc. 85:2149-2156, which is incorporated herein by reference.

 The prepared proteins and fragments thereof can be isolated and purified from the reaction mixture by means of peptide separation, e.g., by extraction, precipitation,
25 electrophoresis, various forms of chromatography, and the like. The receptors of this invention can be obtained in varying degrees of purity depending upon its desired use. Purification can be accomplished by use of the protein purification techniques disclosed herein, see below, or by the use of the antibodies herein described in methods of immunoabsorbant affinity chromatography. This immunoabsorbant affinity
30 chromatography is typically carried out by first linking the antibodies to a solid support and then contacting the linked antibodies with solubilized lysates of appropriate cells, lysates of

other cells expressing the receptor complexes, or lysates or supernatants of cells producing the proteins as a result of DNA techniques, see below.

Generally, the purified protein will be at least about 40% pure, ordinarily at least about 50% pure, usually at least about 60% pure, typically at least about 70% pure, more
5 typically at least about 80% pure, preferable at least about 90% pure and more preferably at least about 95% pure, and in particular embodiments, 97%-99% or more. Purity will usually be on a weight basis, but can also be on a molar basis. Different assays will be applied as appropriate.

10 VI. Antibodies

The term "antibody" or "antibody molecule" as used in this invention includes intact molecules as well as fragments thereof, such as Fab, F(ab')₂, and Fv which are capable of selectively binding the epitopic determinant. These antibody fragments retain some ability to selectively bind with its antigen or receptor and are defined as follows: (1) Fab, the
15 fragment which contains a monovalent antigen-binding fragment of an antibody molecule can be produced by digestion of whole antibody with the enzyme papain to yield an intact light chain and a portion of one heavy chain; (2) Fab', the fragment of an antibody molecule can be obtained by treating whole antibody with pepsin, followed by reduction, to yield an intact light chain and a portion of the heavy chain; two Fab' fragments are obtained per
20 antibody molecule; (3) (Fab')₂, the fragment of the antibody that can be obtained by treating whole antibody with the enzyme pepsin without subsequent reduction; F(ab')₂ is a dimer of two Fab' fragments held together by two disulfide bonds; (4) Fv, defined as a genetically engineered fragment containing the variable region of the light chain and the variable region of the heavy chain expressed as two chains; and (5) Single chain antibody ("SCA"), defined
25 as a genetically engineered molecule containing the variable region of the light chain, the variable region of the heavy chain, linked by a suitable polypeptide linker as a genetically fused single chain molecule.

Methods of making these fragments are known in the art. See, e.g., Harlow and Lane (current edition) Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory,
30 New York. Therefore, the phrase "antibody molecule" in its various forms as used herein contemplates both an intact antibody (immunoglobulin) molecule and an immunologically

active portion of an antibody (immunoglobulin) molecule. Recombinant methods may be applied to make these fragments.

The term "monoclonal antibody" refers to a population of one species of antibody molecule of antigen-specificity. A monoclonal antibody contains only one species of antibody combining site capable of immunoreacting with a particular antigen and thus typically displays a single binding affinity for that antigen. A monoclonal antibody may therefore contain a bispecific antibody molecule having two antibody combining sites, each immunospecific for a different antigen. In one embodiment, the first antibody molecule is affixed to a solid support.

As used in this invention, the term "epitope" means an antigenic determinant on an antigen to which the paratope of an antibody binds. Epitopic determinants usually consist of chemically active surface groupings such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics.

The word "complex" as used herein refers to the product of a specific binding agent-target reaction. An exemplary complex is the combination of two subunits to form a physiological complex of protein subunits, or an immunoreaction product formed by an antibody-antigen reaction.

The term "antigen" refers to a polypeptide or protein that is able to specifically bind to (immunoreact with) an antibody and form an immunoreaction product (immunocomplex). The site on the antigen with which the antibody binds is referred to as an antigenic determinant or epitope, and the binding should be detectable, e.g., 2X, 5X or more above background.

A method of the invention for detection of antibodies that bind to novel epitopes in a sample is performed in vitro, e.g., in immunoassays in which the antibodies can be identified in liquid phase or bound to a solid phase carrier. In various embodiments, the method is performed with a capture antibody bound to a solid support, and/or the capture antibody is a monoclonal antibody molecule. In other instances, the use of tetramer technology may be useful. See, e.g., Kelleher and Rowland-Jones (2000) Curr. Op. Immunol. 12:370-374; Katz (1999) Biomol. Eng. 16:57-65; and Ogg and McMichael (1998) Curr. Op. Immunol. 10:393-396.

Examples of types of immunoassays which can be utilized to detect novel antibodies in a sample, include competitive and non-competitive immunoassays in either a direct or indirect format. Examples of such immunoassays are the radioimmunoassay (RIA) and the sandwich (immunometric) assay. Detection of the antibodies can be done utilizing
5 immunoassays which are run in either the forward, reverse, or simultaneous modes, including competition immunoassays and immunohistochemical assays on physiological samples. Preferably, the method of the invention utilizes a forward immunoassay. Those of skill in the art will know, or can readily discern, other immunoassay formats without undue experimentation.

10 Solid phase-bound antibody molecules are bound by adsorption from an aqueous medium, although other modes of affixation, such as covalent coupling or other well known means of affixation to the solid matrix can be used. Preferably, the first antibody molecule is bound to a support before forming an immunocomplex with antigen, however, the immunocomplex can be formed prior to binding the complex to the solid support.

15 Non-specific protein binding sites on the surface of the solid phase support are preferably blocked. After adsorption of solid phase-bound antibodies, an aqueous solution of a protein free from interference with the assay such as bovine, horse, or other serum albumin that is also free from contamination with the antigen is admixed with the solid phase to adsorb the admixed protein onto the surface of the antibody-containing solid
20 support at protein binding sites on the surface that are not occupied by the antibody molecule.

A typical aqueous protein solution contains about 2-10 weight percent bovine serum albumin in PBS at a pH of about 7-8. The aqueous protein solution-solid support mixture is typically maintained for a time period of at least one hour at a temperature of about 4-37
25 degrees C and the resulting solid phase is thereafter rinsed free of unbound protein.

The first preselected antibody can be bound to many different carriers and used to detect novel epitope binding antibodies in a sample. Examples of well-known carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amyloses, natural and modified celluloses, polyacrylamides, agaroses, and magnetite. The nature of the
30 carrier can be either soluble or insoluble for purposes of the invention. Those skilled in the art will know of other suitable carriers for binding antibodies, or will be able to ascertain such, using routine experimentation.

In addition, if desirable, an antibody for detection in these immunoassays can be detectably labeled in various ways. There are many different labels and methods of labeling known. Examples of the types of labels which can be used in the present invention include, e.g., enzymes, radioisotopes, fluorescent compounds, colloidal metals, chemiluminescent compounds, and bio-luminescent compounds. Many other suitable labels exist for binding to the monoclonal antibodies of the invention

Antibodies which bind to IL-B50 receptor complex can be prepared using an intact polypeptide or fragments containing peptides of interest as the immunizing antigen. The polypeptide or a peptide used to immunize an animal can be derived from translated cDNA or chemical synthesis which can be conjugated to a carrier protein, if desired. Such commonly used carriers which are chemically coupled to the peptide include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid. The coupled peptide is then used to immunize the host animal, e.g., a mouse, a rat, or a rabbit.

If desired, polyclonal or monoclonal antibodies can be further purified, e.g., by binding to and elution from a matrix to which is bound the antigen to which the antibodies were raised. Many techniques are available for purification and/or concentration of polyclonal antibodies, as well as monoclonal antibodies. See, e.g., Coligan, et al. (current ed.) Unit 9, Current Protocols in Immunology, Wiley Interscience.

It is also possible to use the anti-idiotypic antibody technology to produce monoclonal antibodies which mimic an epitope. For example, an anti-idiotypic monoclonal antibody made to a first monoclonal antibody will have a binding domain in the hypervariable region which is the "image" of the epitope bound by the first monoclonal antibody.

The preparation of polyclonal antibodies is well-known to those skilled in the art. See, e.g., Green, et al. "Production of Polyclonal Antisera" pages 1-5 in Manson (ed.) Immunochemical Protocols Humana Press; Production of Polyclonal Antisera in Rabbits, Rats, Mice and Hamsters section 2.4.1 in Coligan, et al. Current Protocols in Immunology.

The preparation of monoclonal antibodies likewise is typically conventional. See, e.g., Kohler and Milstein (1975) Nature 256:495-497; Coligan, et al. Current Protocols sections 2.5.1-2.6.7; and Harlow and Lane (1989). Briefly, monoclonal antibodies can be obtained by injecting mice with a composition comprising an antigen, verifying the presence of antibody production by removing a serum sample, removing the spleen to

obtain B lymphocytes, fusing the B lymphocytes with myeloma cells to produce hybridomas, cloning the hybridomas, selecting positive clones that produce antibodies to the antigen, and isolating the antibodies from the hybridoma cultures. Monoclonal antibodies can be isolated and purified from hybridoma cultures by a variety of well-established techniques. Such isolation techniques include affinity chromatography with Protein-A
5 Sepharose, size-exclusion chromatography, and ion-exchange chromatography. See, e.g., Coligan, et al. Current Protocols sections 2.7.1-2.7.12 and 2.9.1-2.9.3; Barnes, et al. in Methods in Molecular Biology, Humana Press.

Therapeutic applications are conceivable for the antibodies of the present invention.
10 For example, antibodies of the present invention may also be derived from subhuman primate antibody. General techniques for raising therapeutically useful antibodies in baboons may be found, e.g., in Goldenberg, et al. (1991) WO 91/11465; and Losman, et al. (1990) Int. J. Cancer 46:310-314.

Alternatively, a therapeutically useful anti-IL-B50 functional receptor antibody may
15 be derived from a "humanized" monoclonal antibody. Humanized monoclonal antibodies are produced by transferring mouse complementary determining regions from heavy and light variable chains of the mouse immunoglobulin into a human variable domain, and then substituting human residues in the framework regions of the murine counterparts. The use of antibody components derived from humanized monoclonal antibodies obviates potential
20 problems associated with the immunogenicity of murine constant regions. General techniques for cloning murine immunoglobulin variable domains are described, e.g., by Orlandi, et al. (1989) Proc. Nat'l Acad. Sci. USA 86:3833-3837. Techniques for producing humanized monoclonal antibodies are described, e.g., by Jones, et al. (1986) Nature 321:522-525; Riechmann, et al. (1988) Nature 332:323-327; Verhoeven, et al. (1988)
25 Science 239:1534-1536; Carter, et al. (1992) Proc. Nat'l Acad. Sci. USA 89:4285-4289; Sandhu (1992) Crit. Rev. Biotech. 12:437-462; and Singer, et al. (1993) J. Immunol. 150:2844-2857.

Antibodies of the invention also may be derived from human antibody fragments isolated from a combinatorial immunoglobulin library. See, e.g., Barbas, et al. (1991)
30 Methods: A Companion to Methods in Enzymology, vol. 2, page 119; and Winter, et al. (1994) Ann. Rev. Immunol. 12:433-465. Cloning and expression vectors that are useful for

producing a human immunoglobulin phage library can be obtained, e.g., from STRATAGENE Cloning Systems (La Jolla, CA).

In addition, antibodies of the present invention may be derived from a human monoclonal antibody. Such antibodies are obtained from transgenic mice that have been "engineered" to produce specific human antibodies in response to antigenic challenge. In this technique, elements of the human heavy and light chain loci are introduced into strains of mice derived from embryonic stem cell lines that contain targeted disruptions of the endogenous heavy and light chain loci. The transgenic mice can synthesize human antibodies specific for human antigens, and the mice can be used to produce human antibody-secreting hybridomas. Methods for obtaining human antibodies from transgenic mice are described by Green, et al. (1994) Nature Genet. 7:13-21; Lonberg, et al. (1994) Nature 368:856-859; and Taylor, et al. (1994) Int. Immunol. 6:579-591.

Antibody fragments of the present invention can be prepared by proteolytic hydrolysis of the antibody or by expression in *E. coli* of DNA encoding the fragment. Antibody fragments can be obtained by pepsin or papain digestion of whole antibodies by conventional methods. For example, antibody fragments can be produced by enzymatic cleavage of antibodies with pepsin to provide a fragment denoted $F(ab')_2$. This fragment can be further cleaved using a thiol reducing agent, and optionally a blocking group for the sulfhydryl groups resulting from cleavage of disulfide linkages, to produce Fab' monovalent fragments. Alternatively, an enzymatic cleavage using papain produces two monovalent Fab fragments and an Fc fragment directly. These methods are described, e.g., by Goldenberg, U.S. patents No. 4,036,945 and No. 4,331,647, and references contained therein. These patents are hereby incorporated in their entireties by reference including all figures, drawings, and illustrations. See also Nisonhoff, et al. (1960) Arch. Biochem. Biophys. 89:230-244; Porter (1959) Biochem. J. 73:119-127; Edelman, et al. (1967) Methods in Enzymology, vol. 1, Academic Press; and Coligan, et al. Current Protocols, at sections 2.8.1-2.8.10 and 2.10.1-2.10.4.

Other methods of cleaving antibodies, such as separation of heavy chains to form monovalent light-heavy chain fragments, further cleavage of fragments, or other enzymatic, chemical, or genetic techniques may also be used, so long as the fragments bind to the antigen that is recognized by the intact antibody.

For example, Fv fragments comprise an association of V_H and V_L chains. This association may be noncovalent, as described in Inbar, et al. (1972) Proc. Nat'l Acad. Sci. USA 69:2659-2662. Alternatively, the variable chains can be linked by an intermolecular disulfide bond or cross-linked by chemicals such as glutaraldehyde. See, e.g., Sandhu
5 (1992) Crit. Rev. Biotech. 12:437-462. Preferably, the Fv fragments comprise V_H and V_L chains connected by a peptide linker. These single-chain antigen binding proteins (sFv) are prepared by constructing a structural gene comprising DNA sequences encoding the V_H and V_L domains connected by an oligonucleotide. The structural gene is inserted into an expression vector, which is subsequently introduced into a host cell such as E. coli. The
10 recombinant host cells synthesize a single polypeptide chain with a linker peptide bridging the two V domains. Methods for producing sFvs are described, e.g., by Whitlow, et al. (1991) Methods: A Companion to Methods in Enzymology, vol. 2, page 97; Bird, et al. (1988) Science 242:423-426; Ladner, et al., U.S. patent No. 4,946,778; Pack, et al. (1993) Bio/Technology 11:1271-77; and Sandhu (1992) Crit. Rev. Biotech. 12:437-462.

15 Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest. Such genes are prepared, e.g., by using the polymerase chain reaction to synthesize the variable region from RNA of antibody-producing cells. See, e.g., Larrick, et al. (1991) Methods: A
20 Companion to Methods in Enzymology, vol. 2, page 106.

Antibodies can be raised to the various mammalian, e.g., IL-B50 receptor complex and fragments thereof, both in naturally occurring native forms and in their recombinant forms, the difference being that antibodies to the active ligand are more likely to recognize epitopes which are only present in the native conformations. Denatured antigen detection
25 can also be useful in, e.g., Western blot analysis. Anti-idiotypic antibodies are also contemplated, which would be useful as agonists or antagonists of a natural receptor or an antibody.

A number of immunogens may be used to produce antibodies specifically reactive with receptor complex proteins. Recombinant protein is the preferred immunogen for the
30 production of monoclonal or polyclonal antibodies. Naturally occurring protein may also be used either in pure or impure form. Synthetic peptides made using the human IL-B50

receptor protein sequences described herein may also used as an immunogen for the production of antibodies to receptor complexes. Recombinant protein can be expressed in eukaryotic or prokaryotic cells as described herein, and purified as described. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated for subsequent use in immunoassays to measure the protein.

Methods of producing polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized with the mixture. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the antigen of interest. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired. See Harlow and Lane.

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell. Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. The genes encoding the derived antibodies may be subjected to mutagenesis to enhance binding affinity or pharmacokinetic properties. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse, et al. (1989) Science 246:1275-1281.

Antibodies, including binding fragments and single chain versions, against predetermined fragments of the protein can be raised by immunization of animals with conjugates of the fragments with immunogenic proteins. Monoclonal antibodies are prepared from cells secreting the desired antibody. These antibodies can be screened for binding to normal or defective protein, or screened for agonistic or antagonistic activity.

These monoclonal antibodies will usually bind with at least a K_D of about 1 mM, more usually at least about 300 μ M, typically at least about 100 μ M, more typically at least about 30 μ M, preferably at least about 10 μ M, and more preferably at least about 3 μ M or better; including 1 μ M, 300 nM, 100 nM, 30 nM, etc.

5 The antibodies, including antigen binding fragments, of this invention can have significant diagnostic or therapeutic value. They can be potent antagonists that bind to the receptor complex and inhibit binding to the ligand or inhibit the ability of ligand to elicit a biological response. They also can be useful as non-neutralizing antibodies and can be coupled to toxins or radionuclides to bind producing cells. Further, these antibodies can be
10 conjugated to drugs or other therapeutic agents, either directly or indirectly by means of a linker.

 The antibodies of this invention can also be useful in diagnostic applications. As capture or non-neutralizing antibodies, they can bind to the interleukin without inhibiting receptor binding. As neutralizing antibodies, they can be useful in competitive binding
15 assays. They will also be useful in detecting or quantifying functional receptor complex. They may be used as reagents for Western blot analysis, or for immunoprecipitation or immunopurification of the respective protein.

 Protein fragments may be joined to other materials, particularly polypeptides, as fused or covalently joined polypeptides to be used as immunogens. Primate receptor
20 complex and its fragments may be fused or covalently linked to a variety of immunogens, such as keyhole limpet hemocyanin, bovine serum albumin, tetanus toxoid, etc. See Microbiology, Hoeber Medical Division, Harper and Row, 1969; Landsteiner (1962) Specificity of Serological Reactions, Dover Publications, New York; and Williams, et al. (1967) Methods in Immunology and Immunochemistry, Vol. 1, Academic Press, New York;
25 each of which are incorporated herein by reference, for descriptions of methods of preparing polyclonal antisera. A typical method involves hyperimmunization of an animal with an antigen. The blood of the animal is then collected shortly after the repeated immunizations and the gamma globulin is isolated.

 In some instances, it is desirable to prepare monoclonal antibodies from various
30 mammalian hosts, such as mice, rodents, primates, humans, etc. Description of techniques for preparing such monoclonal antibodies may be found in, e.g., Stites, et al. (eds.) Basic

and Clinical Immunology (4th ed.), Lange Medical Publications, Los Altos, CA, and references cited therein; Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH Press; Goding (1986) Monoclonal Antibodies: Principles and Practice (2d ed.) Academic Press, New York; and particularly in Kohler and Milstein (1975) in Nature 256:495-497, which discusses one method of generating monoclonal antibodies. Each of these references is incorporated herein by reference. Summarized briefly, this method involves injecting an animal with an immunogen. The animal is then sacrificed and cells taken from its spleen, which are then fused with myeloma cells. The result is a hybrid cell or "hybridoma" that is capable of reproducing in vitro. The population of hybridomas is then screened to isolate individual clones, each of which secrete a single antibody species to the immunogen. In this manner, the individual antibody species obtained are the products of immortalized and cloned single B cells from the immune animal generated in response to a specific site recognized on the immunogenic substance.

Other suitable techniques involve in vitro exposure of lymphocytes to the antigenic polypeptides or alternatively to selection of libraries of antibodies in phage or similar vectors. See, Huse, et al. (1989) "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," Science 246:1275-1281; and Ward, et al. (1989) Nature 341:544-546, each of which is hereby incorporated herein by reference. The polypeptides and antibodies of the present invention may be used with or without modification, including chimeric or humanized antibodies. Frequently, the polypeptides and antibodies will be labeled by joining, either covalently or non-covalently, a substance which provides for a detectable signal. A wide variety of labels and conjugation techniques are known and are reported extensively in both the scientific and patent literature. Suitable labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent moieties, chemiluminescent moieties, magnetic particles, and the like. Patents, teaching the use of such labels include U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,277,437; 4,275,149; and 4,366,241. Also, recombinant or chimeric immunoglobulins may be produced, see Cabilly, U.S. Patent No. 4,816,567; or made in transgenic mice, see Mendez, et al. (1997) Nature Genetics 15:146-156. These references are incorporated herein by reference.

The antibodies of this invention can also be used for affinity chromatography in isolating primate receptor or cells expressing such. Columns can be prepared where the

antibodies are linked to a solid support, e.g., particles, such as agarose, SEPHADEX, or the like, where a cell lysate may be passed through the column, the column washed, followed by increasing concentrations of a mild denaturant, whereby the purified protein will be released. Conversely, protein may be used to purify antibody.

5 Antibodies may also be used to screen expression libraries for particular expression products. Usually antibodies used in such a procedure will be labeled with a moiety allowing easy detection of presence of antigen by antibody binding.

 Antibodies raised against an IL-B50 receptor will also be used to raise anti-idiotypic antibodies. These will be useful in detecting or diagnosing various immunological
10 conditions related to expression of the protein or cells which express receptors for the protein. They also will be useful as agonists or antagonists of the IL-B50, which may be competitive inhibitors or substitutes for naturally occurring ligands.

Binding composition: Receptor Protein Complex

15 An IL-B50 receptor that specifically binds to or that is specifically immunoreactive with an antibody, e.g., a polyclonal antibody generated against a defined immunogen, e.g., an immunogen consisting of a complex comprising both IL-7R α and R δ 2, is typically determined in an immunoassay. Included within the present invention are those nucleic acid sequences described herein, including functional variants, that encode polypeptides that
20 combine to bind to polyclonal antibodies generated against the prototypical primate IL-B50 receptor, but not the prior known individual components. The immunoassay typically uses a polyclonal antiserum which was raised, e.g., to a complex comprising proteins of SEQ ID NO: 2 and 4. This antiserum is selected to have low crossreactivity against other IL-7
25 receptor family members, preferably from the same species, and to other rodent or similar evolutionarily distant receptors, so that any such crossreactivity is removed by immunoabsorption prior to use in the immunoassay.

 In order to produce antisera for use in an immunoassay, the functional complex is isolated as described herein. For example, recombinant protein may be produced in a mammalian cell line. An appropriate host, e.g., an inbred strain of mice such as Balb/c, is
30 immunized with the complex using a standard adjuvant, such as Freund's adjuvant, and a standard mouse immunization protocol (see Harlow and Lane). Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can

be used as an immunogen. Polyclonal sera are collected, perhaps immunodepleted, and
titered against the immunogen protein in an immunoassay, e.g., a solid phase immunoassay
with the immunogen immobilized on a solid support. Polyclonal antisera with a titer of 10^4
or greater are selected and tested for their cross reactivity against other IL-7 receptor family
5 members, e.g., using a competitive binding immunoassay such as the one described in
Harlow and Lane, supra, at pages 570-573. Preferably at least two IL-7 receptor family
members are used in this determination. These IL-7 receptor family members can be
produced as recombinant proteins and isolated using standard molecular biology and protein
chemistry techniques as described herein.

10 Immunoassays in the competitive binding format can be used for the crossreactivity
determinations. For example, the receptor complex, soluble or membrane associated, can be
immobilized to a solid support. Proteins added to the assay compete with the binding of the
antisera to the immobilized antigen. The ability of the above proteins to compete with the
binding of the antisera to the immobilized complex is compared to the other family
15 members. The percent crossreactivity for the above proteins is calculated, using standard
calculations. Those antisera with less than 10% crossreactivity with each of the proteins
listed above are selected and pooled. The cross-reacting antibodies are then removed from
the pooled antisera by immunoabsorption with the above-listed proteins.

The immunoabsorbed and pooled antisera are then used in a competitive binding
20 immunoassay as described above to compare a second protein complex to the immunogen
protein complex. In order to make this comparison, the two proteins are each assayed at a
wide range of concentrations and the amount of each protein required to inhibit 50% of the
binding of the antisera to the immobilized protein is determined. If the amount of the
second protein required is less than twice the amount of the antigen receptor complex that is
25 required, then the second protein is said to specifically bind to an antibody generated to the
immunogen.

VII. Kits and quantitation

Both naturally occurring and recombinant forms of the receptor complex of this
30 invention are particularly useful in kits and assay methods. For example, these methods
would also be applied to screening for binding activity, e.g., ligands or antagonists for these

receptors. Several methods of automating assays have been developed in recent years so as to permit screening of tens of thousands of compounds per year. See, e.g., a BIOMEK automated workstation, Beckman Instruments, Palo Alto, California, and Fodor, et al. (1991) Science 251:767-773, which is incorporated herein by reference. The latter describes
5 means for testing binding by a plurality of defined polymers synthesized on a solid substrate. The development of suitable assays to screen for an agonist/antagonist or ligand-like proteins can be greatly facilitated by the availability of large amounts of purified, soluble receptor complexes in an active state such as is provided by this invention.

Purified receptor complexes can be attached to substrates for use in the
10 aforementioned screening techniques. However, non-neutralizing antibodies to these receptors can be used as capture antibodies to immobilize the respective receptor complexes on the solid phase, useful, e.g., in diagnostic uses.

This invention also contemplates use of receptor complexes, fragments thereof, peptides, and their fusion products in a variety of diagnostic kits and methods for detecting
15 the presence of the receptor. Alternatively, or additionally, antibodies against the molecules may be incorporated into the kits and methods. Typically the kit will have a compartment containing either a defined peptide or gene segment or a reagent which recognizes one or the other. Typically, recognition reagents, in the case of peptide, would be a ligand or antibody, or in the case of a gene segment, would usually be a hybridization probe.

20 A preferred kit for determining the concentration of, e.g., receptor complex, a sample would typically comprise a labeled compound, e.g., IL-B50 or antibody, having known binding affinity for receptor, a source of receptor (naturally occurring or recombinant) as a positive control, and a means for separating the bound from free labeled compound, e.g., a solid phase for immobilizing the receptor in the test sample.
25 Compartments containing reagents, and instructions, will normally be provided.

Antibodies, including antigen binding fragments, specific for mammalian receptor complex or a peptide fragment, or ligand are useful in diagnostic applications to detect the presence of elevated levels of receptor and/or its fragments. Diagnostic assays may be homogeneous (without a separation step between free reagent and antibody-antigen
30 complex) or heterogeneous (with a separation step). Various commercial assays exist, such as radioimmunoassay (RIA), enzyme-linked immunosorbent assay (ELISA), enzyme immunoassay (EIA), enzyme-multiplied immunoassay technique (EMIT), substrate-labeled

fluorescent immunoassay (SLFIA) and the like. For example, unlabeled antibodies can be employed by using a second antibody which is labeled and which recognizes the antibody to receptor complex or to a particular fragment thereof. These assays have also been extensively discussed in the literature. See, e.g., Harlow and Lane (1988) Antibodies: A Laboratory Manual, CSH., and Coligan (ed. 1991 and periodic supplements) Current Protocols In Immunology Greene/Wiley, New York.

Anti-idiotypic antibodies may have similar use to serve as agonists or antagonists of receptor. These should be useful as therapeutic reagents under appropriate circumstances.

Frequently, the reagents for diagnostic assays are supplied in kits, so as to optimize the sensitivity of the assay. For the subject invention, depending upon the nature of the assay, the protocol, and the label, either labeled or unlabeled antibody, or labeled ligand is provided. This is usually in conjunction with other additives, such as buffers, stabilizers, materials necessary for signal production such as substrates for enzymes, and the like. Preferably, the kit will also contain instructions for proper use and disposal of the contents after use. Typically the kit has compartments for each useful reagent, and will contain instructions for proper use and disposal of reagents. Desirably, the reagents are provided as a dry lyophilized powder, where the reagents may be reconstituted in an aqueous medium having appropriate concentrations for performing the assay.

Many of the aforementioned constituents of the diagnostic assays may be used without modification or may be modified in a variety of ways. For example, labeling may be achieved by covalently or non-covalently joining a moiety which directly or indirectly provides a detectable signal. In any of these assays, a test compound, receptor complex, or antibodies thereto can be labeled either directly or indirectly. Possibilities for direct labeling include label groups: radiolabels such as ^{125}I , enzymes (U.S. Pat. No. 3,645,090) such as peroxidase and alkaline phosphatase, and fluorescent labels (U.S. Pat. No. 3,940,475) capable of monitoring the change in fluorescence intensity, wavelength shift, or fluorescence polarization. Both patents are incorporated herein by reference. Possibilities for indirect labeling include biotinylation of one constituent followed by binding to avidin coupled to one of the above label groups.

There are also numerous methods of separating the bound from the free ligand, or alternatively the bound from the free test compound. The receptor complex can be

immobilized on various matrixes followed by washing. Suitable matrices include plastic such as an ELISA plate, filters, and beads. Methods of immobilizing the receptor to a matrix include, without limitation, direct adhesion to plastic, use of a capture antibody, chemical coupling, and biotin-avidin. The last step in this approach involves the precipitation of antibody/antigen complex by any of several methods including those
5 utilizing, e.g., an organic solvent such as polyethylene glycol or a salt such as ammonium sulfate. Other suitable separation techniques include, without limitation, the fluorescein antibody magnetizable particle method described in Rattle, et al. (1984) Clin. Chem. 30(9):1457-1461, and the double antibody magnetic particle separation as described in U.S.
10 Pat. No. 4,659,678, each of which is incorporated herein by reference.

The methods for linking protein or fragments to various labels have been extensively reported in the literature and do not require detailed discussion here. Many of the techniques involve the use of activated carboxyl groups either through the use of carbodiimide or active esters to form peptide bonds, the formation of thioethers by reaction
15 of a mercapto group with an activated halogen such as chloroacetyl, or an activated olefin such as maleimide, for linkage, or the like. Fusion proteins will also find use in these applications.

Another diagnostic aspect of this invention involves use of oligonucleotide or polynucleotide sequences taken from the sequences of these primate receptor subunits.
20 These sequences can be used as probes for detecting levels of the receptors in patients suspected of having an immunological disorder, or to evaluate polymorphic variation. The preparation of both RNA and DNA nucleotide sequences, the labeling of the sequences, and the preferred size of the sequences has received ample description and discussion in the literature. Normally an oligonucleotide probe should have at least about 14 nucleotides,
25 usually at least about 18 nucleotides, and the polynucleotide probes may be up to several kilobases. Various labels may be employed, most commonly radionuclides, particularly ³²P. However, other techniques may also be employed, such as using biotin modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which may be labeled with a wide variety of labels, such as
30 radionuclides, fluorescers, enzymes, or the like. Alternatively, antibodies may be employed which can recognize specific duplexes, including DNA duplexes, RNA duplexes,

DNA-RNA hybrid duplexes, or DNA-protein duplexes. The antibodies in turn may be labeled and the assay carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected. The use of probes to the novel anti-sense RNA may be carried out in any
5 conventional techniques such as nucleic acid hybridization, plus and minus screening, recombinational probing, hybrid released translation (HRT), and hybrid arrested translation (HART). This also includes amplification techniques such as polymerase chain reaction (PCR).

Diagnostic kits which also test for the qualitative or quantitative presence of other
10 markers are also contemplated. Diagnosis or prognosis may depend on the combination of multiple indications used as markers. Thus, kits may test for combinations of markers. See, e.g., Viallet, et al. (1989) Progress in Growth Factor Res. 1:89-97.

VIII. Therapeutic Utility

15 This invention provides reagents with significant therapeutic value. The receptor (naturally occurring or recombinant), fragments thereof, soluble constructs, and antibodies, along with compounds identified as having binding affinity to the receptor, are useful in the treatment of conditions exhibiting abnormal expression of the cytokine or receptor. Such abnormality will typically be manifested by immunological disorders or otherwise, as
20 described. Additionally, this invention provides therapeutic value in various diseases or disorders associated with abnormal expression or abnormal triggering of response to the interleukin.

In addition, the expression profiles of functional receptor subunits suggests what cells should be responsive to the ligand, e.g., IL-B50. In particular, cells of the lymphoid
25 lineage, e.g., macrophages and dendritic cells, express both subunits in sufficient stoichiometric amounts to form functional complexes. In particular, since the IL-7R α is shared with the IL-7 ligand, it would be expected that much of the signaling pathway will be very similar to IL-7. Thus, much of the biology of the same cells will be closely related. In contrast, the expression of the IL-B50 receptor subunits in different cell types suggests that
30 signaling to those cell types occurs, and will have effects on the physiology effected by such cell types. An antagonist, mutein or antibody, could prove very useful in those situations. See Rich (ed.) Clinical Immunology: Principles and Practice, Mosby.

T helper cells mediate effector functions in infectious, allergic, or autoimmune diseases through production of cytokines. CD4⁺ T cells can be divided into Th1 and Th2 subsets on the basis of their cytokine profile upon antigen stimulation. Evidence has recently been obtained that Th1 and Th2 cells differ in responsiveness and receptor expression. The linkage via TARC expression in response to IL-B50, and production of Th2 type cytokines suggests effect on Th2 responses. The expression profile of the proteins here described indicates that IL-B50 is the ligand for functional receptor complex and, as such, could be important for Th2 effector functions.

Recombinant soluble receptor may be useful as an antagonist, antibodies against the receptor subunits or complex, or IL-B50 antibodies or cytokine muteins can be purified and then administered to a patient. These reagents can be combined for therapeutic use with additional active ingredients, e.g., in conventional pharmaceutically acceptable carriers or diluents, along with physiologically innocuous stabilizers and excipients. These combinations can be sterile, e.g., filtered, and placed into dosage forms as by lyophilization in dosage vials or storage in stabilized aqueous preparations. This invention also contemplates use of antibodies or binding fragments thereof which are not complement binding.

Further analysis can be performed to identify additional molecules involved in the receptor to the cytokine, e.g., additional receptor subunits. Subsequent biological assays can then be utilized to determine if those additional receptor components can affect signal transduction, which can block provide means to further address mechanisms or structures in the signaling pathways. Receptor fragments, e.g., soluble receptor constructs, or ligand muteins can be used as a blocker or antagonist which blocks the activity of IL-B50. Conversely, a compound having intrinsic stimulating activity can activate the receptor and is thus an agonist in that it simulates the activity of IL-B50. This invention further contemplates the therapeutic use of antibodies to the receptor as agonists.

The quantities of reagents necessary for effective therapy will depend upon many different factors, including means of administration, target site, physiological state of the patient, and other medicants administered. Thus, treatment dosages should be titrated to optimize safety and efficacy. Typically, dosages used in vitro may provide useful guidance in the amounts useful for in situ administration of these reagents. Animal testing of effective doses for treatment of particular disorders will provide further predictive indication

of human dosage. Various considerations are described, e.g., in Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, (current ed.), Mack Publishing Co., Easton, Penn.; each of which is hereby incorporated herein by reference. Methods for
5 administration are discussed therein and below, e.g., for oral, intravenous, intraperitoneal, or intramuscular administration, transdermal diffusion, and others. Pharmaceutically acceptable carriers will include water, saline, buffers, and other compounds described, e.g., in the Merck Index, Merck & Co., Rahway, New Jersey. Dosage ranges would ordinarily be expected to be in amounts lower than 1 mM concentrations, typically less than about 10
10 μ M concentrations, usually less than about 100 nM, preferably less than about 10 pM (picomolar), and most preferably less than about 1 fM (femtomolar), with an appropriate carrier. Slow release formulations, or slow release apparatus will often be utilized for continuous administration.

Receptor subunits, complexes of subunits, fragments thereof, and antibodies or their
15 fragments, antagonists, and agonists, or vectors encoding any of the mentioned entities may be administered directly to the host to be treated or, depending on the size of the compounds, it may be desirable to conjugate them to carrier proteins such as ovalbumin or serum albumin prior to their administration. Therapeutic formulations may be administered in many conventional dosage formulations. While it is possible for the active ingredient to
20 be administered alone, it is preferable to present it as a pharmaceutical formulation. Formulations comprise at least one active ingredient, as defined above, together with one or more acceptable carriers thereof. Each carrier must be both pharmaceutically and physiologically acceptable in the sense of being compatible with the other ingredients and not injurious to the patient. Formulations include those suitable for oral, rectal, nasal, or
25 parenteral (including subcutaneous, intramuscular, intravenous and intradermal) administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. See, e.g., Gilman, et al. (eds. 1990) Goodman and Gilman's: The Pharmacological Bases of Therapeutics, 8th Ed., Pergamon Press; and Remington's Pharmaceutical Sciences, 17th ed. (1990), Mack
30 Publishing Co., Easton, Penn.; Avis, et al. (eds. 1993) Pharmaceutical Dosage Forms: Parenteral Medications Dekker, NY; Lieberman, et al. (eds. 1990) Pharmaceutical Dosage

Forms: Tablets Dekker, NY; and Lieberman, et al. (eds. 1990) Pharmaceutical Dosage Forms: Disperse Systems Dekker, NY.

Another therapeutic approach included within the invention involves direct administration of reagents or compositions by any conventional administration techniques (e.g., but not restricted to local injection, inhalation, or administered systemically), to the subject with an appropriate medical disorder. The reagent, formulation, or composition may also be targeted to specific cells or receptors by methods described herein. The actual dosage of reagent, formulation or composition that modulates an inflammatory disorder depends on many factors, including the size and health of an organism. See, e.g., Spilker (1984) Guide to Clinical Studies and Developing Protocols, Raven Press, New York, particularly pages 7-13, 54-60; Spilker (1991) Guide to Clinical Trials, Raven Press, New York, especially pages 93-101; Craig and Stitzel (eds. 1986) Modern Pharmacology 2d ed., Little, Brown, Boston, especially pages 127-33; Speight (ed. 1987) Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3d ed., Williams and Wilkins, Baltimore, pages 50-56; and Tallarida, et al. (1988) Principles in General Pharmacology, Springer-Verlag, New York, pages 18-20; which describes how to determine the appropriate dosage. Generally, doses in the range of between about 0.5 ng/ml and 500 µg/ml inclusive final concentration are administered per day to an adult in a pharmaceutically-acceptable carrier. The therapy of this invention may be combined with or used in association with other therapeutic agents directed to the indicated conditions, particularly agonists or antagonists of other IL-7 family members.

The IL-B50 receptor complex forms the basis for antagonist drug development (e.g., humanized anti-human receptor antibodies). IL-B50 may form an integral part of the lymphoid lineage immune defense. The IL-B50 appears to signal through the STAT3 and STAT5 molecule pathways.

The broad scope of this invention is best understood with reference to the following examples, which are not intended to limit the inventions to the specific embodiments.

The terms and expressions which have been employed are used as terms of description and not of limitation. It is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional

features, modification and variation of the concepts herein disclosed may be used by those of ordinary skill in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

5

EXAMPLES

I. General Methods

Some of the standard methods are described or referenced, e.g., in Maniatis, et al. (1982) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor Press; Sambrook, et al. (1989) Molecular Cloning: A Laboratory Manual, (2d ed.), vols 1-3, CSH Press, NY; Ausubel, et al., Biology, Greene Publishing Associates, Brooklyn, NY; or Ausubel, et al. (1987 and Supplements) Current Protocols in Molecular Biology, Greene/Wiley, New York. Methods for protein purification include such methods as ammonium sulfate precipitation, column chromatography, electrophoresis, centrifugation, crystallization, and others. See, e.g., Ausubel, et al. (1987 and periodic supplements); Deutscher (1990) "Guide to Protein Purification" in Meth. Enzymol., vol. 182, and other volumes in this series; and manufacturer's literature on use of protein purification products, e.g., Pharmacia, Piscataway, N.J., or Bio-Rad, Richmond, CA. Combination with recombinant techniques allow fusion to appropriate segments, e.g., to a FLAG sequence or an equivalent which can be fused via a protease-removable sequence. See, e.g., Hochuli (1989) Chemische Industrie 12:69-70; Hochuli (1990) "Purification of Recombinant Proteins with Metal Chelate Absorbent" in Setlow (ed.) Genetic Engineering, Principle and Methods 12:87-98, Plenum Press, N.Y.; and Crowe, et al. (1992) QIAexpress: The High Level Expression & Protein Purification System QIAGEN, Inc., Chatsworth, CA.

Computer sequence analysis is performed, e.g., using available software programs, including those from the GCG (U. Wisconsin) and GenBank sources. Public sequence databases were also used, e.g., from GenBank and others.

Many techniques applicable to the IL-10 receptor may be applied to IL-7R α and/or R δ 2, as described, e.g., in U.S. Patent No. 5,985,828, which is incorporated herein by reference for all purposes.

30

II. Amplification of receptor fragments by PCR

Cloning of human receptors is performed by standard procedures. Various methods of amplifying target sequences, such as the polymerase chain reaction, can also be used to prepare DNA encoding these receptor proteins or polypeptides. Polymerase chain reaction (PCR) technology is used to amplify such nucleic acid sequences directly from mRNA, from cDNA, and from genomic libraries or cDNA libraries. This allows for diagnostic methods which allow determination of polymorphic or populational variety, many of which might affect physiology or function of the resulting gene product. Such may allow predictive information to be gathered to diagnose disease or predict therapeutic response.

In PCR techniques, oligonucleotide primers complementary to two 5' regions in the DNA region to be amplified are synthesized. The polymerase chain reaction is then carried out using the two primers. See Innis et al. (current eds.) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA. Primers can be selected to amplify the entire regions encoding full-length receptor proteins or to amplify smaller DNA segments as desired. Once such regions are PCR-amplified, they can be sequenced and oligonucleotide probes can be prepared from sequence obtained using standard techniques.

Oligonucleotides for use as probes are chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Carruthers (1983) Tetrahedron Lett. 22(20): 1859-1862, or using an automated synthesizer, as described in Needham-VanDevanter et al. (1984) Nucleic Acids Res. 12: 6159-6168. Purification of oligonucleotides is performed, e.g., by native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson and Regnier (1983) J. Chrom. 255: 137-149. The sequence of the synthetic oligonucleotide can be verified using the chemical degradation method of Maxam and Gilbert in Grossman and Moldave (eds. 1980) Methods in Enzymology 65:499-560 Academic Press, New York.

The peptide segments, along with comparison to homologous genes, can also be used to produce appropriate oligonucleotides to screen a library. The genetic code can be used to select appropriate oligonucleotides useful as probes for screening. In combination with polymerase chain reaction (PCR) techniques, synthetic oligonucleotides will be useful in selecting desired clones from a library.

Complementary sequences will also be used as probes or primers. Based upon identification of the likely amino terminus, other peptides should be particularly useful, e.g.,

coupled with anchored vector or poly-A complementary PCR techniques or with complementary DNA of other peptides.

To identify a homologous receptor protein, degenerate oligonucleotides are designed. The primers are used for polymerase chain reactions on, e.g., genomic DNA
5 followed by subcloning the PCR products using restriction sites placed at the 5' ends of the primers, picking individual *E. coli* colonies carrying these subcloned inserts, and using a combination of random sequencing and hybridization analysis to eliminate other known family members.

Subsequently, PCR products are gel-purified, digested with appropriate restriction
10 enzymes, gel-purified again, and subcloned in the Bluescript vector (Stratagene, San Diego, CA). Bacterial colonies carrying individual subclones are picked into 96 well microtiter plates, and multiple replicas are prepared by plating the cells onto nitrocellulose. The replicate filters are hybridized to probes representing known members of the IL-7 receptor family, and DNA is prepared from non-hybridizing colonies for sequence analysis.

Two appropriate forward and reverse primers are selected using the sequences
15 supplied herein (see Figures 1 and 2) and common knowledge. See, e.g., Innis, et al. (current eds.) PCR Protocols: A Guide to Methods and Applications Academic Press, San Diego, CA; and Dieffenbach and Dveksler (current eds.) PCR Primer: A Laboratory Manual Cold Spring Harbor Press, CSH, NY. RT-PCR is used on an appropriate mRNA sample
20 selected for the presence of message to produce a cDNA, e.g., a monocyte or macrophage cell sample.

Full length clones may be isolated by hybridization of cDNA libraries from appropriate tissues pre-selected by PCR signal.

As is commonly known, PCR primers are typically designed to contain at least 15
25 nucleotides, e.g., 15-30 nucleotides. The design of specific primers containing 21 nucleotides, e.g., that code for the appropriate polypeptides containing at least 4 amino acids from the IL7R α or R δ 2 sequences are described as follows. Other PCR primers designed to amplify other receptor polypeptide fragments will be designed in a similar fashion, e.g., mutagenesis primers. Preferably, most or all of the nucleotides in such a primer encode
30 conserved amino acids, e.g., amino acid residues of SEQ ID NO: 2 or 4. For example, primers containing at least 40% IL-7R α or R δ 2 conserved amino acids can be used. Once

appropriate amino acids are selected as templates against which primer sequences are to be designed, the primers can be synthesized using, e.g., standard chemical methods. Due to the degeneracy of the genetic code and the bias of preferred species variants, such primers should be designed to include appropriate degenerate sequences, as can be readily
5 determined using common knowledge.

III. Tissue distribution of IL-7R α or R δ 2

Message for the gene encoding IL-7R α has been detected in a human cDNA library,
10 in dendritic cells and monocytes. Message for R δ 2 has been detected in dendritic cells, monocytes, NK, and some T cells. Both are found in certain DC and certain monocyte samples, indicating that the components of functional receptors are on those cells.

Southern Analysis: DNA (5 μ g) from a primary amplified cDNA library is digested with appropriate restriction enzymes to release the inserts, run on a 1% agarose gel and
15 transferred to a nylon membrane (Schleicher and Schuell, Keene, NH).

Samples for human mRNA isolation could include: peripheral blood mononuclear cells (monocytes, T cells, NK cells, granulocytes, B cells), resting (T100); peripheral blood mononuclear cells, activated with anti-CD3 for 2, 6, 12 h pooled (T101); T cell, TH0 clone Mot 72, resting (T102); T cell, TH0 clone Mot 72, activated with anti-CD28 and anti-CD3
20 for 3, 6, 12 h pooled (T103); T cell, TH0 clone Mot 72, anergic treated with specific peptide for 2, 7, 12 h pooled (T104); T cell, TH1 clone HY06, resting (T107); T cell, TH1 clone HY06, activated with anti-CD28 and anti-CD3 for 3, 6, 12 h pooled (T108); T cell, TH1 clone HY06, anergic treated with specific peptide for 2, 6, 12 h pooled (T109); T cell, TH2 clone HY935, resting (T110); T cell, TH2 clone HY935, activated with anti-CD28 and anti-
25 CD3 for 2, 7, 12 h pooled (T111); T cells CD4+CD45RO- T cells polarized 27 days in anti-CD28, IL-4, and anti IFN- γ , TH2 polarized, activated with anti-CD3 and anti-CD28 4 h (T116); T cell tumor lines Jurkat and Hut78, resting (T117); T cell clones, pooled AD130.2, Tc783.12, Tc783.13, Tc783.58, Tc782.69, resting (T118); T cell random $\gamma\delta$ T cell clones, resting (T119); Splenocytes, resting (B100); Splenocytes, activated with anti-CD40 and IL-
30 4 (B101); B cell EBV lines pooled WT49, RSB, JY, CVIR, 721.221, RM3, HSY, resting (B102); B cell line JY, activated with PMA and ionomycin for 1, 6 h pooled (B103); NK 20

clones pooled, resting (K100); NK 20 clones pooled, activated with PMA and ionomycin for 6 h (K101); NKL clone, derived from peripheral blood of LGL leukemia patient, IL-2 treated (K106); NK cytotoxic clone 640-A30-1, resting (K107); hematopoietic precursor line TF1, activated with PMA and ionomycin for 1, 6 h pooled (C100); U937 premonocytic line, resting (M100); U937 premonocytic line, activated with PMA and ionomycin for 1, 6 h pooled (M101); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 1, 2, 6, 12, 24 h pooled (M102); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 1, 2, 6, 12, 24 h pooled (M103); elutriated monocytes, activated with LPS, IFN γ , anti-IL-10 for 4, 16 h pooled (M106); elutriated monocytes, activated with LPS, IFN γ , IL-10 for 4, 16 h pooled (M107); elutriated monocytes, activated LPS for 1 h (M108); elutriated monocytes, activated LPS for 6 h (M109); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, resting (D101); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 1 hr (D102); DC 70% CD1a+, from CD34+ GM-CSF, TNF α 12 days, activated with PMA and ionomycin for 6 hr (D103); DC 95% CD1a+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D104); DC 95% CD14+, ex CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin 1, 6 hr pooled (D105); DC CD1a+ CD86+, from CD34+ GM-CSF, TNF α 12 days FACS sorted, activated with PMA and ionomycin for 1, 6 h pooled (D106); DC from monocytes GM-CSF, IL-4 5 days, resting (D107); DC from monocytes GM-CSF, IL-4 5 days, resting (D108); DC from monocytes GM-CSF, IL-4 5 days, activated LPS 4, 16 h pooled (D109); DC from monocytes GM-CSF, IL-4 5 days, activated TNF α , monocyte supe for 4, 16 h pooled (D110); leiomyoma L11 benign tumor (X101); normal myometrium M5 (O115); malignant leiomyosarcoma GS1 (X103); lung fibroblast sarcoma line MRC5, activated with PMA and ionomycin for 1, 6 h pooled (C101); kidney epithelial carcinoma cell line CHA, activated with PMA and ionomycin for 1, 6 h pooled (C102); kidney fetal 28 wk male (O100); lung fetal 28 wk male (O101); liver fetal 28 wk male (O102); heart fetal 28 wk male (O103); brain fetal 28 wk male (O104); gallbladder fetal 28 wk male (O106); small intestine fetal 28 wk male (O107); adipose tissue fetal 28 wk male (O108); ovary fetal 25 wk female (O109); uterus fetal 25 wk female (O110); testes fetal 28 wk male (O111); spleen fetal 28 wk male (O112); adult placenta 28 wk (O113); and tonsil inflamed, from 12 year old (X100).

IV. Production of mammalian receptor protein

Typically, co-expression will be useful, and constructs can be made to produce soluble receptors. This may be effected by a polycistronic construct in appropriate cells, or
5 by construction of vectors which coexpress both receptor subunits together. Alternatively, fusion constructs can be produced, e.g., to generate antibodies.

An appropriate construct is engineered for expression, e.g., in *E. coli*. For example, a mouse IGIF pGex plasmid is constructed and transformed into *E. coli*. Freshly transformed cells are grown in LB medium containing 50 µg/ml ampicillin and induced with
10 IPTG (Sigma, St. Louis, MO). After overnight induction, the bacteria are harvested and the pellets containing receptors are isolated. The pellets are homogenized in TE buffer (50 mM Tris-base pH 8.0, 10 mM EDTA and 2 mM pefabloc) in 2 liters. This material is passed through a microfluidizer (Microfluidics, Newton, MA) three times. The fluidized supernatant is spun down on a Sorvall GS-3 rotor for 1 h at 13,000 rpm. The resulting
15 supernatant containing the receptor is filtered and passed over a glutathione-SEPHAROSE column equilibrated in 50 mM Tris-base pH 8.0. The fractions containing the receptor protein are pooled and cleaved with thrombin (Enzyme Research Laboratories, Inc., South Bend, IN). The cleaved pool is then passed over a Q-SEPHAROSE column equilibrated in 50 mM Tris-base. Fractions containing receptor are pooled and diluted in cold distilled
20 H₂O, to lower the conductivity, and passed back over a fresh Q-SEPHAROSE column. Fractions containing receptor are pooled, aliquoted, and stored in the -70 degrees C freezer.

Comparison of the CD spectrum with functional receptor may suggest that the protein is correctly folded. See Hazuda, et al. (1969) *J. Biol. Chem.* 264:1689-1693.

Protein expression and purification of human receptor: Adenoviral vectors
25 containing full-length human receptors are constructed by PCR and used to transfect, e.g., Q293 packaging cells. Produced viruses are subsequently purified, with all procedures according to manufacturer's protocols (Invitrogen). Receptor proteins are prepared, e.g., from 5 x 10⁸ infected Q293 cells (adenoviruses at 10 MOI) which are subsequently incubated for 5 days in a cell factory in a total volume of 1 l of serum-free CMF-1 medium
30 (Gibco BRL). Culture medium is dialyzed (Spectra/Por membrane tubing, MW cut off: 6-8 kD) against 50 mM Tris-HCl, pH 8.0, 1 mM EDTA, and subsequently passed over Hitrap Q

sepharose and Heparin columns. The flow-through, containing the receptor proteins, is sterile-filtered and concentrated approximately 70 times using an Amicon 8400 ultrafiltration cell with a 10 kD MW cut off membrane. The samples are dialyzed against PBS, and the protein content quantified by PAGE and Coomassie Blue staining using lysozyme as a standard. Protein identities are confirmed, e.g., by N-terminal sequencing. Identically treated Q293 cells infected with adenovirus encoding green fluorescent protein provide a mock control. Endotoxin levels are determined using the Limulus Amebocyte Lysate assay (BioWhittaker) and were less than 4 EU/ml. Protein samples are stored at 4 degrees C.

Expression vectors: For mammalian expression, vectors encoding full-length human receptor are constructed by inserting PCR-generated cDNA fragments into pME18S. Kitamura, et al. (1991) Proc. Nat'l Acad. Sci. USA 88:5082-5086.

V. Preparation of antibodies specific for receptor complex

Inbred Balb/c mice are immunized intraperitoneally with recombinant forms of the protein, e.g., purified soluble receptor-FLAG or stable transfected NIH-3T3 cells. Animals are boosted at appropriate time points with protein or cells, with or without additional adjuvant, to further stimulate antibody production. Serum is collected, or hybridomas produced with harvested spleens.

Alternatively, Balb/c mice are immunized with cells transformed with the gene or fragments thereof, either endogenous or exogenous cells, or with isolated membranes enriched for expression of the antigen. Serum is collected at the appropriate time, typically after numerous further administrations. Various gene therapy techniques may be useful, e.g., in producing protein in situ, for generating an immune response.

Monoclonal antibodies may be made. For example, splenocytes are fused with an appropriate fusion partner and hybridomas are selected in growth medium by standard procedures. Hybridoma supernatants are screened for the presence of antibodies which bind to the desired receptor, e.g., by ELISA or other assay. Antibodies which specifically recognize receptor or complex may also be selected or prepared.

In another method, synthetic peptides or purified protein are presented to an immune system to generate monoclonal or polyclonal antibodies. See, e.g., Coligan (1991) Current Protocols in Immunology Wiley/Greene; and Harlow and Lane (1989) Antibodies: A

Laboratory Manual Cold Spring Harbor Press. In appropriate situations, the binding reagent is either labeled as described above, e.g., fluorescence or otherwise, or immobilized to a substrate for panning methods. Nucleic acids may also be introduced into cells in an animal to produce the antigen, which serves to elicit an immune response. See, e.g., Wang, et al. (1993) Proc. Nat'l. Acad. Sci. 90:4156-4160; Barry, et al. (1994) BioTechniques 16:616-619; and Xiang, et al. (1995) Immunity 2: 129-135.

VI. Biological Activity of IL-B50 and Complexes with IL-7R α and R δ 2

A number of experiments were conducted in order to assess the signaling receptor complex for IL-B50, as well as the biological function of IL-B50. In the following examples, the term "human TSLP" or "hTSLP" is used interchangeably with the term IL-B50. Additionally, the term "human TSLPR" or "hTSLPR" is used interchangeably with the term R δ 2. Materials and methods used in the following experiments were as follows.

Cell lines. Human 293T epithelial cells were maintained in Dulbecco's modified Eagle's medium (DMEM) (Life Technologies Inc.) supplemented with 10% fetal calf serum (FCS) (LTBMC). The Pro-B cell line Ba/F3 was maintained in RPMI 1640 medium (Life Technologies Inc.) supplemented with 10% fetal calf serum and 10 ng/ml of mouse IL-3. QBI-293A human embryonic kidney cells used for adenovirus expression were grown in CMF-1 medium (CellWorks, San Diego, CA). BOSC23 cells were maintained in DMEM-10% FCS and guanine phosphoribosyltransferase (GPT) selection reagents (Specialty Media). The cells were transferred to Dulbecco's modified Eagle's medium-10% FCS without GPT selection reagents 2 days before transfection.

Adenovirus expression of human TSLP (IL-B50) and purification of the recombinant protein. The mature coding region of human TSLP (residues 1-131 of Figure 3) was fused to the signal sequence of mouse SLAM (Bates, et al. (1999) J. Immunol. 163:1973) and inserted into a modified version of transfer vector pQB1-AdCMV5-GFP (Quantum Biotechnologies Inc.) by PCR. Recombinant adenovirus was produced as described in Quantum applications manual 24AL98. Recombinant virus was used to infect 5×10^8 cells in

1 L CMF-1 with culture in a Nunc Cell Factory (Nalge Nunc Int., Naperville, IL) for 3 days. The culture medium was clarified by centrifugation, dialyzed and filtered prior to application to a 5 ml Q-Sepharose column. The Q-Sepharose flow-through, which contained human TSLP, was loaded onto a 5 ml HiTrap Heparin (Pharmacia, Uppsala, Sweden) column at 5 ml/min. The column was washed with 50 mM Tris-HCl pH 8.0, 1 mM EDTA, and eluted with a gradient from 0-2.5 M NaCl in 50 mM Tris-HCl pH 8.0, 1 mM EDTA. The peak fractions were concentrated, dialyzed against PBS and quantitated by SDS-PAGE and Coomassie staining using lysozyme as a standard. A similar procedure was followed to prepare mouse TSLP.

10

Ba/F3 retroviral-mediated gene transfer and proliferation assays. Human IL-7R cDNA and human TSLPR cDNA were cloned by PCR in the retroviral vectors pMX and its derivative pMX-puro to give pMX-hIL-7R α and pMX-puro-TSLPR, respectively (Kitamura, T. (1998) Int. J. Hematol. 67:351). The BOSC23 packaging cell line was transiently transfected with retrovirus constructs using Fugene 6 (GIBCO BRL) according to the manufacturer protocol. Retrovirus containing supernatants were collected after two days. Ba/F3 cells were infected with retroviral supernatants for 48 hr on petri dishes coated with 40 μ g/ml recombinant fibronectin fragments (Retronectin, Takara). After 48 hr puromycin (1 μ g/ml) was added to those cells infected with virus obtained from pMX-puro constructs. The efficiency of infection of Ba/F3 cells was over 90% as assessed by parallel infection with the test construct pMXI-EGFP encoding the enhanced green fluorescent protein (EGFP). Proliferation assays using Ba/F cells were as previously described (Ho, et al. (1993) Proc. Natl. Acad. Sci. USA 90:11267). Cells were washed three times with RPMI media and plated at a density of 5000 cells/well. Cells were grown with serial threefold dilutions of mouse IL-3, human and mouse TSLP, or human IL-7 (all starting concentrations 225 ng/ml). After 36 hr at 37 degrees C Alamar Blue \otimes REDOX indicator (Trek Diagnostic systems) was added to a final concentration of 10% (vol/vol) to each well. Cells were allowed to grow for 5-8 more hours after which plates were measured with a fluorometer.

20
25
30

Quantitation of mRNA expression. cDNA libraries from various tissues and cellular sources were prepared as described previously (Bolin, et al. (1997) J. Neurosci. 17:5493)

and used as templates for Taqman-PCR analyses. cDNAs (50 ng per reaction) were analyzed for the expression of hTSLP, hTSLPR and hIL7R α genes by the Fluorogenic 5'-nuclease PCR assay (Holland, et al. (1991) Proc. Natl. Acad. Sci. USA 88:7276), using an ABI Prism 7700 Sequence Detection System (Perkin Elmer, Foster City, CA). Reactions were incubated for 2 min at 50 degrees C, denatured for 10 min at 95 degrees C and subjected to 40 two-step amplification cycles with annealing/extension at 60 degrees C for 1 min followed by denaturation at 95 degrees C for 15 sec. The amplicons used for hTSLP, hTSLPR and IL-7R α covered bp 246-315, bp 263-335 and bp 519-596, respectively (numbering starts at start codon), and were analyzed with FAM-labeled probes. Values were expressed as fg/50 ng total cDNA. Primers and probes for human chemokine and chemokine receptors were obtained from Perkin-Elmer as PreDeveloped Assay Reagents (PDAR's). Chemokine and chemokine receptor expression was adjusted for the amount of 18SrRNA and compared to the control (calibrator) samples using the comparative C_T method (Fehniger, et al. (1999) J. Immunol. 162:4511). Samples were measured in duplicate. 18SrRNA levels were determined under primer limited conditions in multiplex reactions as recommended using a Vic labelled probe (Perkin Elmer, Foster City, CA).

Cell isolation and culture. Peripheral Blood Mononuclear Cells (PBMC) were purified from buffy coats of healthy volunteers (Stanford Blood Bank, Palo Alto, CA) by centrifugation over Ficoll. Human monocytes were isolated from PBMC by negative depletion using anti-CD2 (Leu-5A), anti-CD3 (Leu-4), anti-CD8 (Leu 2a), anti-CD19 (Leu-12), anti-CD20 (Leu-16), anti-CD56 (Leu-19), (BD, San Jose CA), anti-CD67 (IOM 67) (Immunotech, Westbrook, ME) and anti-glycophorin A (10F7MN, ATCC, Rockville, MD) mAbs and sheep anti-mouse IgG coupled magnetic beads (Dyna, Oslo, Norway) as described previously (Koppelman, et al. (1997) Immunity 7:861). Monocytes were cultured in RPMI + 10% FCS at a density of 10^6 cells/ml in the presence or absence of IL-7 (50 ng/ml) and/or hTSLP (50 ng/ml) for 24 hrs and culture supernatants and cells were harvested for quantitation of cytokine production or gene expression analyses. Human CD11c+ dendritic cells (DC) were isolated from PBMC as previously described (Kadowaki, et al. (2000) J. Exp. Med. 192:219). Briefly, PBMC were incubated with anti-CD3, anti-CD14, anti-CD19, anti-CD56 mAbs, depleted from lineage+ cells using magnetic beads

(Dyna) and CD11c+ Lineage-blood DC were subsequently isolated by cell sorting to reach a purity of more than 99%. Freshly sorted cells were cultured in RPMI1640 containing 10% FCS at 5×10^4 /100 μ l in flat-bottom 96-well half-area culture plates or at 1×10^5 /200 μ l in flat-bottom 96-well plates, with or without IL-B50 (15 ng/ml).

5

TARC Elisa. The production of TARC/CCL17 in culture supernatants was determined by chemokine specific elisa using MAB364 as capture reagent and BAF364 as detection reagent (R&D Systems, Minneapolis MN). The sensitivity of the assay was 50 pg/ml.

10

DC viability and flow cytometric analysis. After 24 hours of culture, DC were harvested and resuspended in an EDTA-containing medium to dissociate the clusters. Viable DC were first counted using trypan blue exclusion of dead cells. Remaining cells were stained with a variety of mouse anti-human FITC-conjugated monoclonal antibodies (mAb) including anti-HLA-DR (Becton Dickinson), anti-CD40, anti-CD80 and CD86 (all from Pharmingen) or an Ig-G1 isotype control (Becton Dickinson), and were analyzed with a FACScan[®] flow cytometer (Becton Dickinson). Dead cells were excluded based on side and forward scatter characteristics.

15

T cell proliferation assay. Naïve CD4+/CD45RA+ T cells were isolated from adult blood buffy coats by negative depletion of cells expressing CD14, CD19, CD56, CD8, CD45RO, HLA-DR and glycophorin A using magnetic beads (Dyna). More than 95% of the purified cells had the CD4+CD45RA+ naïve T cell phenotype. CD11c+ DC were washed twice to remove any cytokine and co-cultured with 5×10^4 allogeneic naïve CD4+ T cells in round-bottom 96-well culture plates at increasing DC/T cell ratios. All co-cultures were carried out in triplicate. DC alone and T cells alone were used as controls. After 5 days, cells were pulsed with 1 μ Ci ³H-Thymidine (Amersham) for 16 hours before harvesting and counting radioactivity.

20

25

30

Stat3 and Stat5 activation assays. Stable Ba/F3 transfectant cells ($\sim 2.5 \times 10^7$ cells) were starved for 4-6 hours, and then stimulated at 10^6 cells/ml for 15 min with either 10

ng/ml of mIL-3 or 30 ng/ml of hTSLP. After stimulation, cells were harvested and incubated for 15 min at 4 degrees C in lysis buffer containing 50 mM Tris-HCL pH 7.5, 300 mM NaCl, 2 mM EDTA, 0.875% Brij 97, 0.125% NP40, 10 mg/ml aprotinin, 10 mg/ml leupeptin, 1 mM PMSF, 1 mM Na₃VO₄, and 1 mM NaF. Cell lysates were clarified by
5 centrifugation at 12,000 X g for 15 min, and supernatants were subjected to 8% SDS-PAGE. Proteins were electrotransferred onto nylon membranes (Immobilon-P, Millipore, Bradford, MA) and detected by Western Blot analysis using rabbit Abs against anti-phospho Stat3 and anti-phospho Stat5 (New England Biolabs) or anti-Stat3 and anti-Stat5 (Santa Cruz Biotechnology), followed by mouse anti-rabbit Ig HRP. Immunoreactive bands were
10 visualized with enhanced chemiluminescence (ECL) (SuperSignal West Dura Extended Duration Substrate, Pierce, Rockford, IL) on ECL film (Kodak). For reprobing, blots were stripped with 200 mM glycine, 1% SDS, pH 2.5 for 30 min at 65 degrees C.

VI.A. IL-B50 signals via IL-7R α and R δ 2

15 The cytokine human IL-B50 has as closest homologs human and mouse IL-7 and the recently described mouse TSLP (Sims, et al. (2000) *J. Expt'l Med.* 192:671-680). In mouse, both IL-7 and TSLP function as T- and B-cell growth and differentiation factors. The signaling receptor complexes for mouse IL-7 and mouse TSLP consist of two subunits, respectively mouse IL-7R α and mouse R γ c (common receptor for IL-2, IL-4, IL-7, IL-9, and IL-15) for mouse IL-7, and mouse IL-7R α and mouse TSLPR (R δ 1) for the mTSLP
20 ligand (Park, et al. (2000) *J. Expt'l Med.* 192:659-670).

In an attempt to identify the signaling receptor complex for human IL-B50, Ba/F3 cells were co-transfected with expression constructs for human IL-7R α and an orphan human cytokine receptor known as R δ 2, a subunit related to R γ c and mTSLPR (R δ 1), using
25 the methods described above. Any functional relationship between the mTSLPR subunit R δ 1 and the human R δ 2 had been unclear. Co-transfected Ba/F3 cells showed a proliferative response in the presence of hIL-B50, but not with hIL-7 or medium. Ba/F3 cells transfected with either hIL-7R α or hR δ 2 alone did not show a proliferative response with hIL-B50. Additionally, no cross-reactivity between mTSLP and the hTSLP receptor
30 complex was observed. These findings establish the signaling complex for human IL-B50 as consisting of hIL-7R α and hR δ 2.

The corresponding activation status of Stat5 and Stat3 was also measured in the various BaF3 cell populations. Both Stat5 and Stat3 were phosphorylated upon addition of hTSLP, but only when both hTSLPR and hIL-7R α was present

5 **VI.B. Expression Analysis of IL-B50, hIL-7R α and hR δ 2**

In order to identify target cells capable of responding to IL-B50888888, a large panel of cDNA libraries was analyzed for the simultaneous expression of both hIL-7R α and hR δ 2, using quantitative PCR. Results of the expression analysis, conducted as described in materials and methods, are presented in Figures 4A-4E. In particular, expression analysis of
10 the two receptor subunits indicated that they were co-expressed primarily in activated dendritic cells, monocytes, and T cells (see, Figures 4A-4D) indicating that these cell types respond to human IL-B50. As shown in Figure 4E, IL-B50 was expressed in various tissue types, with high expression in the human lung.

15 **VI.C. Human IL-B50 Induces Chemokine Expression on Freshly Isolated Monocyte Population and CD11c⁺ blood DC**

The spectrum of biological activities induced by IL-B50 was investigated based on the overlapping expression patterns of IL-B50 receptor components. cDNA was prepared from human monocytes cultured for 24 hrs in the presence of IL-B50 or IL-7, and the expression
20 of 38 human chemokines and 20 human chemokine receptors were analyzed by quantitative "real time" PCR. Interestingly, IL-B50 (TSLP) and IL-7 influenced the expression of distinct sets of chemokines (Table 1), but did not affect the expression of chemokine receptors. IL-B50 enhanced the expression of TARC/CCL17, DC-CK1/PARC/CCL18, MDC/CCL22, and MIP3 β /CL19. IL-7 also enhanced expression of TARC/CCL17,
25 MDC/CCL22, and MIP3 β /CCL19 but in addition, enhanced expression of IL-8/CXCL8, CTAPIII/CXCL7, ENA78/CXCL5, and GROabg/CXCL123 and decreased the expression levels of IP-10/CXCL10, I-TACK/CXCL11, SDF1/CXCL12, MCP2/CCL8 and MCP4/CCL13 (Table 1).

Additionally, the ability of IL-B50 to stimulate DCs to produce mRNAs for various
30 cytokines and chemokines was compared with that of GM-CSF, IL-7, CD40-ligand (CD40L) and medium alone as a control, as follows. Purified CD11c⁺ DCs were cultured

for 15-17 hours with IL-B50 (15 ng/ml), GM-CSF (100 ng/ml), IL-7 (50 ng/ml), CD40-ligand transfected L-cells (1 L-cell/4 DC) or medium alone. Total RNA was extracted and studied using real time quantitative PCR as described above. As shown in Figures 14A and 14C, IL-B50 did not stimulate human DCs to produce mRNA for IL-1 α , IL-1 β , IL-6, IL-12p40, TNF- α , MCP-1, MCP-4, Rantes and MIG, but did stimulate human DCs to produce mRNA for the chemokines TARC, MDC and MIP3- β (Figure 14B).

The induction of TARC protein by IL-B50 on monocyte and dendritic cell populations was confirmed by ELISA. The level of TARC production by CD11c+ dendritic cells was at least ten-fold higher than that of monocytes (Figure 5).

The induction of IL12p75 protein by IL-B50 was also examined. To do so, purified CD11c+ DCs were culture for 24 hours with IL-B50 (15 ng/ml), IL-7 (50 ng/ml), CD40-ligand-transfected L-cells (1 L-cell/4 DC), bacterial lipopolysaccharide (LPS: 1mg/ml) or medium alone. Culture supernatant was harvested and bioactive IL12p75 was measured using a high-sensitivity ELISA kit. As shown in Figure 15, IL-B50 did not stimulate human DCs to produce IL12p75 protein.

Table 1. Effects of IL-B50 and IL-7 on Chemokine Expression*

Chemokine	Media	IL-B50	IL-7
CCL1	40.0	1.0	1.4
CCL2	24.8	1.0	1.5
CCL3	31.5	0.7	2.0
CCL4	28.9	1.0	1.6
CCL5	30.4	1.1	0.6
CCL7	31.3	0.7	1.4
CCL8	30.2	0.8	0.2
CCL11	40.0	1.6	1.4
CCL13	37.3	1.6	0.1
CCL14	40.0	1.3	1.3
CCL15	40.0	1.1	1.3
CCL16	40.0	2.2	7.0

CCL17	39.8	195.4	20.1
CCL18	35.8	2.9	1.7
CCL19	36.7	8.5	8.3
CCL20	40.0	1.2	1.2
CCL21	40.0	1.0	1.0
CCL22	34.3	8.8	3.0
CCL24	29.3	2.0	1.2
CCL25	40.0	1.1	1.1
CCL26	38.9	1.1	2.5
CCL27	40.0	0.8	1.2
CCL28	40.0	1.0	1.0
CXCL1-3	28.7	1.0	4.4
CXCL4	27.9	1.2	1.9
CXCL5	28.7	1.1	8.0
CXCL6	40.0	1.5	1.3
CXCL7	28.7	1.1	2.0
CXCL8	27.3	1.4	8.5
CXCL9	34.9	0.6	0.7
CXCL10	29.7	0.6	0.1
CXCL11	32.9	0.7	0.0
CXCL12	33.1	0.6	0.2
CXCL13	35.4	0.2	0.8
CXCL14	39.7	1.0	1.0
XCL1	40.0	0.9	3,8
CX3CL1	40.0	1.4	1.3

*Human monocytes were cultured in the absence or presence of IL-B50 (50 ng/ml) or IL-7 (50 ng/ml) for 18 h, and expression of chemokine genes was determined by quantitative PCR. Results are expressed as 1) C_T values of nonactivated samples and 2) fold difference relative to the calibrator sample (media).

VL.D. IL-B50 Activates CD11c+ Dendritic Cells.

Freshly purified immature CD11c+ blood DC are known to spontaneously mature in culture. As shown in Figure 6A, loose and irregular clumps in the DC culture were evident after 24 hrs in medium alone. In the presence of IL-B50, this maturation process was dramatically enhanced. DC in culture formed tight and round clumps with fine dendrites

visible at the periphery of each clump (Figure 6B). The IL-B50-induced maturation was confirmed by analyzing the surface phenotype of DC using flow cytometry. Whereas IL-B50 slightly upregulated the expression of HLA-DR and CD86, it strongly induced the costimulatory molecules CD40 and CD80 (Figure 7). This maturation process was accompanied by an increased viability of the DC. Additionally, IL-B50 was more potent than CD40-ligand (CD40L) and IL-7 in upregulating CD40 and CD80 (Figure 8). A titration of IL-B50 using log dilutions of the cytokine showed that both the effect on survival and the induction of costimulatory molecules on DC was maximal at 15 ng/ml and above, and still significant at concentrations as low as 15 pg/ml.

The T cell stimulatory capacity of CD11+ DC, cultured 24 hrs in medium alone or in the presence of IL-B50, was analyzed. DCs were cocultured with 5×10^4 naïve CD4+CD45RA+ allogeneic T cells at increasing DC/T cell ratios. As assessed by ^3H -thymidine incorporation at day 5 of the coculture, DC cultured with IL-B50 induced up to 10-fold stronger naïve T cell proliferation as compared to DC cultured in medium (Figure 9).

Additionally, the T cell stimulatory capacity of CD11+ DC, cultured with LPS, medium alone, IL-7, CD40-ligand and IL-B50, was compared by determining ^3H -thymidine incorporation. As shown in Figure 10, IL-B50-activated DCs were more potent than DCs activated with CD40-ligand, IL-7 and LPS in inducing proliferation of allogeneic naïve CD4 T cells.

DCs, cultured in medium alone, IL-B50, CD40-ligand, IL-7 and LPS, were cocultured with naïve CD4 T cells. After 6 days of coculture, CD4 T cells were restimulated for 24 hours with anti-CD3 and anti-CD28 and the culture supernatants were analyzed by ELISA to quantify the cytokine production by T cells. As shown in Figures 11A-11E, IL-B50-activated human DCs primed naïve CD4 T cells to produce IL-4, IL-13 and TNF- α , but inhibited production of IL-10 and IFN- γ .

DCs were cultured for 24 hours in medium alone, IL-B50, IL-7, CD40-ligand or LPS, to prime purified naïve CD8 T cells over 6 days in coculture. IL-B50-activated DCs strongly induced expansion of allogeneic naïve CD8 T cells (Figure 12), as well as the expression of perforin (Figure 13).

The data herein indicate, among other things, that human IL-B50 is a novel hematopoietic cytokine most closely related to IL-7. It represents a human ortholog of mTSLP. The human IL-B50 signaling makes use of the combination of hIL-7R α and hR δ 2, which together form a novel hematopoietic cytokine receptor. Both receptor subunits are notably present on macrophages and dendritic cells, indicating functional effects of the cytokine on those cell types, and mediating functions provided by those cell types. The human IL-B50 also promotes the phosphorylation of STAT3 and STAT5 transcription factors.

Therapeutic uses of IL-B50 are apparent. For example, SCID patients with mutations in IL-7R α are T-cell deficient. Since IL-B50 uses IL-7R α , IL-B50 share signaling pathway components, and may play a significant role in human T-cell differentiation and may enhance T-cell recovery in circumstances of T-cell depletion.

Likewise, IL-B50 antagonists are useful. The antagonists take various forms such as ligand muteins, antibodies to ligand, and antibodies to receptor, e.g., which block ligand binding. Since the hIL-B50 likely plays a role in the development of T- and B-cell lymphomas, then blocking IL-B50 signaling, either at the ligand or at its receptor components, is useful in treatment of some of these lymphomas.

Herein, based upon the binding studies, hIL-B50 receptor subunit mapping identified IL-7R α and novel human receptor R δ 2 (a close relative of human γ c or IL-2R γ) as signaling receptor complex (co-expression in Ba/F3 cells delivers a proliferative signal in response to IL-B50). Receptor expression profiles indicate both IL-7R α and R δ 2 are primarily expressed on dendritic cells, though they are both also expressed in monocytes. The dendritic cell expression indicates a role for the cytokine in maturation of cells or pathways important in antigen presentation, indicating use of the cytokine for expansion, e.g., *ex vivo*, of antigen presenting cells.

All citations herein are incorporated herein by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference including all figures and drawings.

Many modifications and variations of this invention, as will be apparent to one of ordinary skill in the art can be made to adapt to a particular situation, material, composition

of matter, process, process step or steps, to preserve the objective, spirit and scope of the invention. All such modifications are intended to be within the scope of the claims appended hereto without departing from the spirit and scope of the invention. The specific embodiments described herein are offered by way of example only, and the invention is to
5 be limited by the terms of the appended claims, along with the full scope of equivalents to which such claims are entitled; and the invention is not to be limited by the specific embodiments that have been presented herein by way of example.

WHAT IS CLAIMED IS:

1. A method of producing a ligand:receptor complex, comprising contacting:
 - a) a substantially pure or recombinant mammalian IL-B50 with a receptor
5 comprising the IL-7R α or the R δ 2 subunit;
 - b) a mammalian IL-B50 with a receptor comprising a substantially pure or recombinant IL-7R α subunit; or
 - c) a mammalian IL-B50 with a receptor comprising a substantially pure or recombinant R δ 2 subunit;10 thereby allowing said complex to form.
2. The method of Claim 1, wherein:
 - a) said mammalian IL-B50 is primate IL-B50;
 - b) said complex formation results in signal transduction, STAT activation, or
15 TARC, PARC or MDC expression;
 - c) said receptor is on a cell;
 - d) said receptor comprises both IL-7R α and R δ 2 subunit;
 - e) said complex formation results in a physiological change in the cell expressing said receptor;
 - 20 f) said contacting is in combination with a proliferative agent, cytokine, or chemokine; or
 - g) said contacting allows quantitative detection of said ligand.
3. The method of Claim 2, wherein said receptor is on a hematopoietic cell,
25 including a lymphoid lineage cell, myeloid, or dendritic cell.
4. A method of modulating physiology or development of an IL-7R α or R δ 2 expressing cell comprising contacting said cell to an exogenous agonist or antagonist of a mammalian IL-B50.

5. The method of Claim 4, wherein:

A) said antagonist is:

- 1) an antibody which neutralizes said mammalian IL-B50;
- 2) a mutein of said IL-B50; or
- 3) an antibody which binds to IL-7R α or R δ 2 or a complex comprising both;

B) said physiology is selected from:

- 1) proliferation;
- 2) lymphoid or dendritic lineage cell development;
- 3) antigen presentation; or
- 4) production of inflammatory mediators, including cytokines, chemokines,
or adhesion molecules; or

C) said cell is a hematopoietic cell.

6. The method of Claim 4, wherein:

- a) said antagonist is an antibody and said physiology is hematopoietic cell proliferation;
- b) said agonist is IL-B50 and said physiology is hematopoietic cell differentiation;
or
- c) said physiology is antigen presentation.

7. The method of Claim 4, wherein said modulating is blocking, and said physiology is lymphoid lineage cell proliferation.

8. A method of modulating a signal to a cell mediated by IL-B50 comprising contacting said cell to an administered agonist or antagonist of IL-B50.

9. The method of Claim 8, wherein said modulating is inhibiting, and said signal is a proliferation signal.

10. The method of Claim 9, wherein:
- a) said antagonist is a neutralizing antibody to IL-7R α or the R δ 2 subunit or a complex comprising said subunits;
 - b) said agonist or antagonist is administered in combination with another antagonist or agonist of IL-B50; or
 - c) said agonist or antagonist is administered in combination with a growth factor, cytokine, chemokine, or immune adjuvant.
11. The method of Claim 9, wherein said contacting is with another anti-proliferative agent or treatment.
12. A method of selectively labeling a population of cells, said method comprising contacting said cells with an antibody which binds:
- a) IL-7R α ;
 - b) R δ 2; or
 - c) a complex comprising one of said subunits;
- thereby resulting in the identification of cells expressing said subunit or complex.
13. The method of Claim 12, wherein:
- a) said contacting results in modulation of STAT activation;
 - b) said labeling allows purification of IL-7R α or R δ 2 subunit expressing cells; or
 - c) said labeling allows depletion of IL-7R α or R δ 2 subunit expressing cells.
14. A population of cells made by the method of Claim 13.
15. The population of Claim 14, which are prepared by Fluorescent Activated Cell Sorting.
16. A method of testing a compound for ability to affect receptor-ligand interaction, said method comprising comparing the interaction of a receptor complex

comprising IL-7R α and/or R δ 2 subunit with IL-B50 in the presence and absence of said compound.

17. The method of Claim 16, wherein said compound is an antibody which binds
5 one of:
- a) IL-7R α ;
 - b) R δ 2 subunit;
 - c) a receptor comprising IL-7R and/or R δ 2; or
 - d) IL-B50.
18. An isolated or recombinant protein complex comprising:
- a) at least 15 contiguous amino acid residues of SEQ ID NO: 2 and at least 15
contiguous amino acid residues of SEQ ID NO: 4;
 - b) at least two distinct segments of at least 8 contiguous amino acid residues of SEQ
15 ID NO: 2 and at least two distinct segments of at least 8 contiguous amino
acid residues of SEQ ID NO: 4; or
 - c) at least one segment at least 21 contiguous nucleotides of SEQ ID NO: 1 and at
least one segment at least 21 contiguous nucleotides of SEQ ID NO: 3.
19. The complex of Claim 18, wherein:
- a) one of said segments of SEQ ID NO: 2 is from the extracellular portion of said
sequence;
 - b) one of said segments of SEQ ID NO: 4 is from the extracellular portion of said
sequence; or
 - c) said polypeptide comprises the mature SEQ ID NO: 2 and the mature SEQ ID
25 NO: 4 sequences.
20. An isolated or recombinant polynucleotide encoding said complex of Claim
19, wherein:
- a) said polynucleotide comprises a deoxyribonucleotide;
 - b) said polynucleotide comprises a ribonucleotide; or
- 30

c) at least one of said segments is operably linked to a promoter.

21. A binding compound comprising an antigen binding portion from an antibody which binds with selectivity to a polypeptide comprising at least 12 contiguous amino acid residues of SEQ ID NO: 2 and at least 12 contiguous amino acid residues of
5 SEQ ID NO: 4; said polypeptide encoded by a nucleotide of Claim 20.

Nucleotide and amino acid sequences (see SEQ ID NO: 1 and 2) of a primate,
e.g., human, IL-7R α ; predicted signal cleavage site indicated.

```

ctctctctct atctctctca ga atg aca att cta ggt aca act ttt ggc atg      52
Met Thr Ile Leu Gly Thr Thr Phe Gly Met
-20
-15
gtt ttt tct tta ctt caa gtc gtt tct gga gaa agt ggc tat gct caa      100
Val Phe Ser Leu Leu Gln Val Val Ser Gly Gln Ser Gly Tyr Ala Gln
-10
-5
aat gga gac ttg gaa gat gca gaa ctg gat gac tac tca ttc tca tgc      148
Asn Gly Asp Leu Gln Asp Ala Gln Leu Asp Asp Tyr Ser Phe Ser Cys
10
15
tat agc cag ttg gaa gtg aat gga tcg cag cat tca ctg acc tgt gct      196
Tyr Ser Gln Leu Gln Val Asn Gly Ser Gln His Ser Leu Thr Cys Ala
25
30
ttt gag gac cca gat gtc aac acc acc aat ctg gaa ttg gaa ata tgt      244
Phe Gln Asp Pro Asp Val Asn Thr Thr Asn Leu Gln Phe Gln Ile Cys
40
45
ggg gcc ctg gtc gtc gta aag tgc ctg aat ttc agg aaa cta caa gag      292
Gly Ala Leu Val Gln Val Lys Cys Leu Asn Phe Arg Lys Leu Gln Gln
55
60
65
70

```

FIG. 1A

```

ata tat ttc atc gag aca aag aaa ttc tta ctg att gga aag agc aat      340
Ile Tyr Phe Ile Glu Thr Lys Lys Phe Leu Leu Ile Gly Lys Ser Asn      85
75
ata tgt gtg aag gtt gga gaa aag agt cta acc tgc aaa aaa ata gac      388
Ile Cys Val Lys Val Gly Glu Lys Ser Leu Thr Cys Lys Lys Ile Asp      100
90
cta acc act ata gtt aaa cct gag gct cct ttt gac ctg agt gtc atc      436
Leu Thr Thr Ile Val Lys Pro Glu Ala Pro Phe Asp Leu Ser Val Ile      115
105
tat cgg gaa gga gcc aat gac ttt gtg gtg aca ttt aat aca tca cac      484
Tyr Arg Glu Gly Ala Asn Asp Phe Val Val Thr Phe Asn Thr Ser His      130
125
ttg caa aag aag tat gta aaa gtt tta atg cat gat gta gct tac cgc      532
Leu Glu Lys Lys Tyr Val Lys Val Leu Met His Asp Val Ala Tyr Arg      150
140
cag gaa aag gat gaa aac aaa tgg acg cat gtg aat tta tcc agc aca      580
Gln Glu Lys Asp Glu Asn Lys Trp Thr His Val Asn Leu Ser Ser Thr      165
155

```

FIG. 1B

```

aag ctg aca ctc ctg cag aga aag ctc caa ccg gca gca atg tat gag      628
Lys Leu Thr Leu Leu Gln Arg Lys Leu Gln Pro Ala Ala Met Tyr Glu
170
att aaa gtt cga tcc atc cct gat cac tat ttt aaa ggc ttc tgg agt      676
Ile Lys Val Arg Ser Ile Pro Asp His Tyr Phe Lys Gly Phe Trp Ser
185
gaa tgg agt cca agt tat tac ttc aga act cca gag atc aat aat agc      724
Glu Trp Ser Pro Ser Tyr Tyr Phe Arg Thr Pro Glu Ile Asn Asn Ser
200
tca ggg gag atg gat cct atc tta cta acc atc agc att ttg agt ttt      772
Ser Gly Glu Met Asp Pro Ile Leu Leu Thr Ile Ser Ile Leu Ser Phe
215
ttc tct gtc gct ctg ttg gtc atc ttg gcc tgt gtg tta tgg aaa aaa      820
Phe Ser Val Ala Leu Leu Val Ile Leu Ala Cys Val Leu Trp Lys Lys
235
agg att aag cct atc gta tgg ccc agt ctc ccc gat cat aag aag act      868
Arg Ile Lys Pro Ile Val Trp Pro Ser Leu Pro Asp His Lys Lys Thr
250
ctg gaa cat ctt tgt aag aaa cca aga aaa aat tta aat gtg agt ttc      916
Leu Glu His Leu Cys Lys Lys Pro Arg Lys Asn Leu Asn Val Ser Phe
265

```

FIG. 1C

aat cct gaa agt ttc ctg gac tgc cag att cat agg gtg gat gac att	964
Asn Pro Glu Ser Phe Leu Asp Cys Gln Ile His Arg Val Asp Asp Ile	
280	285
caa gct aga gat gaa gtg gaa ggt ttt ctg caa gat acg ttt cct cag	1012
Gln Ala Arg Asp Glu Val Glu Gly Phe Leu Gln Asp Thr Phe Pro Gln	
295	300
caa cta gaa gaa tct gag aag cag agg ctt gga ggg gat gtg cag agc	1060
Gln Leu Glu Glu Ser Glu Lys Gln Arg Leu Gly Gly Asp Val Gln Ser	
315	320
ccc aac tgc cca tct gag gat gta gtc gtc act cca gaa agc ttt gga	1108
Pro Asn Cys Pro Ser Glu Asp Val Val Val Thr Pro Glu Ser Phe Gly	
330	335
aga gat tca tcc ctg tca tgc ctg gct ggg aat gtc agt gca tgt gac	1156
Arg Asp Ser Ser Leu Thr Cys Leu Ala Gly Asn Val Ser Ala Cys Asp	
345	350
gcc cct att ctg tcc tct tcc agg tcc cta gac tgc agg gag agt ggc	1204
Ala Pro Ile Leu Ser Ser Ser Arg Ser Leu Asp Cys Arg Glu Ser Gly	
360	365
aag aat ggg cct cat gtg tac cag gac ctg ctg ctt agc ctt ggg act	1252
Lys Asn Gly Pro His Val Tyr Gln Asp Leu Leu Leu Ser Leu Gly Thr	
375	380
	385

FIG. 1D

```

aca aac agc acg ctg ccc cct cca ttt tct ctc caa tct gga atc ctg      1300
Thr Asn Ser Thr Leu Pro Pro Pro Phe Ser Leu Gln Ser Gly Ile Leu      405
395

aca ttg aac cca gtt gct cag ggt cag ccc att ctt act tcc ctg gga      1348
Thr Leu Asn Pro Val Ala Gln Gly Gln Pro Ile Leu Thr Ser Leu Gly      420
410

tca aat caa gaa gaa gca tat gtc acc atg tcc agc ttc tac caa aac      1396
Ser Asn Gln Gln Gln Ala Tyr Val Thr Met Ser Ser Phe Tyr Gln Asn      435
425

cag tgaagtgtaa gaaaccacaga ctgaacttac cgtgagcgcac aaagatgatt      1449
Gln

taaaagggaa gtctagagt cctagtctcc ctcaacagcac agagaagaca aaatagcaa      1509

aaccccaacta cacagtctgc aagattctga aacattgctt tgaccactct tcctgagttc      1569

agtggcactc aacatgagtc aagagcatcc tgcttctacc atgtgattt ggtcacaagg      1629

tttaaggtga cccaatgatt cagctattt      1658

```

FIG. 1E

Nucleotide and amino acid sequences (see SEQ ID NO: 3 and 4) of a primate,
e.g., human, R52; predicted signal cleavage site indicated.

```

cggcagcagag gc atg ggg cgg ctg gtt ctg ctg tgg gga gct gcc gtc ttt 51
Met Gly Arg Leu Val Leu Leu Trp Gly Ala Ala Val Phe
-20 -15 -10

ctg ctg gga ggc tgg atg gct ttg ggg caa gga gga gca gca gaa gga 99
Leu Leu Gly Gly Trp Met Ala Leu Gly Gln Gly Gly Ala Ala Gln Gly
-5 -1 5

gta cag att cag atc atc tac ttc aat tta gaa acc gtg cag gtg aca 147
Val Gln Ile Gln Ile Ile Tyr Phe Asn Leu Glu Thr Val Gln Val Thr
10 15 20

tgg aat gcc agc aaa tac tcc agg acc aac ctg act ttc cac tac aga 195
Trp Asn Ala Ser Lys Tyr Ser Arg Thr Asn Leu Thr Phe His Tyr Arg
25 30 35

ttc aac ggt gat gag gcc tat gac cag tgc acc aac tac ctt ctc cag 243
Phe Asn Gly Asp Glu Ala Tyr Asp Gln Cys Thr Asn Tyr Leu Leu Gln
40 45 50 55

gaa ggt cac act tcg ggg tgc ctc cta gac gca gag cag cga gac gac 291
Glu Gly His Thr Ser Gly Cys Leu Leu Asp Ala Glu Gln Arg Asp Asp
60 65 70

```

FIG. 2A

```

att ctc tat ttc tcc atc agg aat ggg acg cac ccc gtt ttc acc gca      339
Ile Leu Tyr Phe Ser Ile Arg Asn Gly Thr His Pro Val Phe Thr Ala
75
75
agt cgc tgg atg gtt tat tac ctg aaa ccc agt tcc ccg aag cac gtg      387
Ser Arg Trp Met Val Tyr Tyr Leu Lys Pro Ser Ser Pro Lys His Val
90
90
aga ttt tcg tgg cat cag gat gca gtg acg gtg acg gtt tct gac ctg      435
Arg Phe Ser Trp His Gln Asp Ala Val Thr Val Thr Cys Ser Asp Leu
105
105
tcc tac ggg gat ctc ctc tat gag gtt cag tac cgg agc ccc ttc gac      483
Ser Tyr Gly Asp Leu Leu Tyr Glu Val Gln Tyr Arg Ser Pro Phe Asp
120
125
acc gag tgg cag tcc aaa cag gaa aat acc tgc aac gtc acc ata gaa      531
Thr Glu Trp Gln Ser Lys Gln Glu Asn Thr Cys Asn Val Thr Ile Glu
140
145
ggc ttg gat gcc gag aag tgt tac tct ttc tgg gtc agg gtg aag gct      579
Gly Leu Asp Ala Glu Lys Cys Tyr Ser Phe Trp Val Arg Val Lys Ala
155
160
atg gag gat gta tat ggg cca gac aca tac cca agc gac tgg tca gag      627
Met Glu Asp Val Tyr Gly Pro Asp Thr Tyr Pro Ser Asp Trp Ser Glu
170
175
180

```

FIG. 2B

```

gtg aca tgc tgg cag aga ggc gag att cgg gat gcc tgt gca gag aca 675
Val Thr Cys Trp Gln Arg Gly Glu Ile Arg Asp Ala Cys Ala Glu Thr
185 190 195
cca acg cct ccc aaa cca aag ctg tcc aaa ttt att tta att tcc agc 723
Pro Thr Pro Pro Lys Pro Lys Leu Ser Lys Phe Ile Leu Ile Ser Ser
200 205 210
ctg gcc atc ctt ctg atg atg tct ctc ctc ctt ctg tct tta tgg aaa 771
Leu Ala Ile Leu Leu Met Val Ser Leu Leu Leu Ser Leu Trp Lys
220 225 230
tta tgg aga gtg aag aag ttt ctc att ccc agc gtg cca gac ccg aaa 819
Leu Trp Arg Val Lys Lys Phe Leu Ile Pro Ser Val Pro Asp Pro Lys
235 240 245
tcc atc ttc ccc ggg ctc ttt gag ata cac caa ggg aac ttc cag gag 867
Ser Ile Phe Pro Gly Leu Phe Glu Ile His Gln Gly Asn Phe Gln Glu
250 255 260
tgg atc aca gac acc cag aac gtg gcc cac ctc cac aag atg gca ggt 915
Trp Ile Thr Asp Thr Gln Asn Val Ala His Leu His Lys Met Ala Gly
265 270 275
gca gag caa gaa agt ggc ccc gag gag ccc ctg gta gtc cag ttg gcc 963
Ala Glu Gln Glu Ser Gly Pro Glu Glu Pro Leu Val Val Gln Leu Ala
280 285 290

```

FIG. 2C

aag act gaa gcc gag tct ccc agg atg ctg gac cca cag acc gag gag	1011
Lys Thr Glu Ala Glu Ser Pro Arg Met Leu Asp Pro Gln Thr Glu Glu	
300	310
aaa gag gcc tct ggg gga tcc ctc cag ctt ccc cac cag ccc ctc caa	1059
Lys Glu Ala Ser Gly Gly Ser Leu Gln Leu Pro His Gln Pro Leu Gln	
315	320
ggc ggt gat gtg gtc aca atc ggg ggc ttc acc ttg atg atg aat gac	1107
Gly Gly Asp Val Val Thr Ile Gly Gly Phe Thr Phe Val Met Asn Asp	
330	335
cgc tcc tac gtg gcg ttg tgatggacac accactgtca aagtcacgt	1155
Arg Ser Tyr Val Ala Leu	
345	

FIG. 2D

```
caagatccac gttgacattt aaagacagag gggactgtcc cggggactcc acaccaccat 1215
ggatgggaaag tctccacgcc aatgatgta ggaactagag actctgaaga cccagcctca 1275
ccgcctaatag cggccactgc cctgctaact ttccccaca tgagtctctg tgtcaaagg 1335
cttgatggca gatgggagcc aattgctcca ggaatttac tccaagttcc tttcgtgcc 1395
tgaacglttgt cacataaacc ccaaggcagc acgtccaanaa tgctgtaaaa ccatcttccc 1455
actctgtgag tccccagttc cgtccatgta cctgttccat agcattgga tctcggaaga 1515
ttttttgtct gtttgagac tccaaccac ctctaccct ac 1557
```

FIG. 2E

```

MEPEALLVLSVSEFRKIFILQLVGLVLTVD 30
FTNCDFEIKIKAYLSTISKDLITYMSGTKS 60
TEFNNTVSCSNRPHCLTEIQSLTFNPAGC 90
ASIAKEMFAMKTKAIAIWCPGYSETQINA 120
TQAMKKRRKRKVTTNKCLEQVSQLOGLMR 150
FNRPLLKQO

```

FIG. 3A


```


1 agtgtgaac tggggtggaa tgggtgtcca cgtatgttcc ctttgcctt
51 actatatgtt ctgtcagttt cttcaggaa aatcttcac ttacaacttg
101 tagggctggt gtttaacttac gacttcacca actgtgactt tgagaagatt
151 aaagcagcct atctcagtac tatttctaaa gacctgatta catatatgag
201 tgggaccaaa agtaccgagt tcaacaacac cgtctctgt agcaatcggc
251 cacattgcct tactgaatc cagagcctaa cctcaatcc caccgccgc
301 tgcgcgtcgc tcgccaaaaga aatggttcgcc atgaaaacta agctgcctt
351 agctatctgg tggccaggtt attcggaac tcagataaat gctactcagg
401 caatgaagaa gaggagaaaa aggaagtca caaccaataa atgtctgaa
451 caagtgtcac aattacaag attgtggcgt cgcttcaatc gacccttact
501 gaaacaacag taaccatct ttattatgt catatttcac agcaccaaaa
ta

```


FIG. 3B


12/35





14/35








FIG. 4E




FIG. 5




FIG. 6A




FIG. 6B

19/35




FIG. 7




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9

24/35





FIG. 10

IL-4**FIG. 11A**




FIG. 11B





FIG. 11C

28/35

IL-10**FIG. 11D**




FIG. 11E




FIG. 12

31/35

Med-DCs

CD8

CD40L-DC

IL-50-DCs




FIG. 13




FIG. 14A




FIG. 14B

34/35




FIG. 14C




FIG. 15

SEQUENCE LISTING

SEQ ID NO: 1 provides primate IL-7R α nucleotide sequence.
 SEQ ID NO: 2 provides primate IL-7R α polypeptide sequence.
 SEQ ID NO: 3 provides primate R δ 2 nucleotide sequence.
 SEQ ID NO: 4 provides primate R δ 2 polypeptide sequence.
 SEQ ID NO: 5 provides primate IL-B50 nucleotide sequence.
 SEQ ID NO: 6 provides primate IL-B50 polypeptide sequence.

<110> Schering Corporation

<120> Mammalian Cytokines; Receptors; Related Reagents and
 Methods

<130> DX01341 PCT

<140>

<141>

<150> US 60/298,268

<151> 2001-06-14

<150> US 60/247,218

<151> 2000-11-10

<160> 6

<170> PatentIn Ver. 2.0

<210> 1

<211> 1658

<212> DNA

<213> Unknown

<220>

<223> Description of Unknown Organism: primate; surmised
 human

<220>

<221> CDS

<222> (23)..(1399)

<220>

<221> mat_peptide

<222> (83)..(1399)

<400> 1
 ctctctctct atctctctca ga atg aca att cta ggt aca act ttt ggc atg 52
 Met Thr Ile Leu Gly Thr Thr Phe Gly Met
 -20 -15

gtt ttt tct tta ctt caa gtc gtt tct gga gaa agt ggc tat gct caa 100
 Val Phe Ser Leu Leu Gln Val Val Ser Gly Glu Ser Gly Tyr Ala Gln
 -10 -5 -1 1 5

aat gga gac ttg gaa gat gca gaa ctg gat gac tac tca ttc tca tgc 148
 Asn Gly Asp Leu Glu Asp Ala Glu Leu Asp Asp Tyr Ser Phe Ser Cys
 10 15 20

tat agc cag ttg gaa gtg aat gga tcg cag cat tca ctg acc tgt gct 196

Tyr Ser Gln Leu Glu Val Asn Gly Ser Gln His Ser Leu Thr Cys Ala
 25 30 35
 ttt gag gac cca gat gtc aac acc acc aat ctg gaa ttt gaa ata tgt 244
 Phe Glu Asp Pro Asp Val Asn Thr Thr Asn Leu Glu Phe Glu Ile Cys
 40 45 50
 ggg gcc ctc gtg gag gta aag tgc ctg aat ttc agg aaa cta caa gag 292
 Gly Ala Leu Val Glu Val Lys Cys Leu Asn Phe Arg Lys Leu Gln Glu
 55 60 65 70
 ata tat ttc atc gag aca aag aaa ttc tta ctg att gga aag agc aat 340
 Ile Tyr Phe Ile Glu Thr Lys Lys Phe Leu Leu Ile Gly Lys Ser Asn
 75 80 85
 ata tgt gtg aag gtt gga gaa aag agt cta acc tgc aaa aaa ata gac 388
 Ile Cys Val Lys Val Gly Glu Lys Ser Leu Thr Cys Lys Lys Ile Asp
 90 95 100
 cta acc act ata gtt aaa cct gag gct cct ttt gac ctg agt gtc atc 436
 Leu Thr Thr Ile Val Lys Pro Glu Ala Pro Phe Asp Leu Ser Val Ile
 105 110 115
 tat cgg gaa gga gcc aat gac ttt gtg gtg aca ttt aat aca tca cac 484
 Tyr Arg Glu Gly Ala Asn Asp Phe Val Val Thr Phe Asn Thr Ser His
 120 125 130
 ttg caa aag aag tat gta aaa gtt tta atg cat gat gta gct tac cgc 532
 Leu Gln Lys Lys Tyr Val Lys Val Leu Met His Asp Val Ala Tyr Arg
 135 140 145 150
 cag gaa aag gat gaa aac aaa tgg acg cat gtg aat tta tcc agc aca 580
 Gln Glu Lys Asp Glu Asn Lys Trp Thr His Val Asn Leu Ser Ser Thr
 155 160 165
 aag ctg aca ctc ctg cag aga aag ctc caa ccg gca gca atg tat gag 628
 Lys Leu Thr Leu Leu Gln Arg Lys Leu Gln Pro Ala Ala Met Tyr Glu
 170 175 180
 att aaa gtt cga tcc atc cct gat cac tat ttt aaa ggc ttc tgg agt 676
 Ile Lys Val Arg Ser Ile Pro Asp His Tyr Phe Lys Gly Phe Trp Ser
 185 190 195
 gaa tgg agt cca agt tat tac ttc aga act cca gag atc aat aat agc 724
 Glu Trp Ser Pro Ser Tyr Tyr Phe Arg Thr Pro Glu Ile Asn Asn Ser
 200 205 210
 tca ggg gag atg gat cct atc tta cta acc atc agc att ttg agt ttt 772
 Ser Gly Glu Met Asp Pro Ile Leu Leu Thr Ile Ser Ile Leu Ser Phe
 215 220 225 230
 ttc tct gtc gct ctg ttg gtc atc ttg gcc tgt gtg tta tgg aaa aaa 820
 Phe Ser Val Ala Leu Leu Val Ile Leu Ala Cys Val Leu Trp Lys Lys
 235 240 245
 agg att aag cct atc gta tgg ccc agt ctc ccc gat cat aag aag act 868
 Arg Ile Lys Pro Ile Val Trp Pro Ser Leu Pro Asp His Lys Lys Thr
 250 255 260

ctg gaa cat ctt tgt aag aaa cca aga aaa aat tta aat gtg agt ttc 916
 Leu Glu His Leu Cys Lys Lys Pro Arg Lys Asn Leu Asn Val Ser Phe
 265 270 275

aat cct gaa agt ttc ctg gac tgc cag att cat agg gtg gat gac att 964
 Asn Pro Glu Ser Phe Leu Asp Cys Gln Ile His Arg Val Asp Asp Ile
 280 285 290

caa gct aga gat gaa gtg gaa ggt ttt ctg caa gat acg ttt cct cag 1012
 Gln Ala Arg Asp Glu Val Glu Gly Phe Leu Gln Asp Thr Phe Pro Gln
 295 300 305 310

caa cta gaa gaa tct gag aag cag agg ctt gga ggg gat gtg cag agc 1060
 Gln Leu Glu Glu Ser Glu Lys Gln Arg Leu Gly Gly Asp Val Gln Ser
 315 320 325

ccc aac tgc cca tct gag gat gta gtc gtc act cca gaa agc ttt gga 1108
 Pro Asn Cys Pro Ser Glu Asp Val Val Thr Pro Glu Ser Phe Gly
 330 335 340

aga gat tca tcc ctc aca tgc ctg gct ggg aat gtc agt gca tgt gac 1156
 Arg Asp Ser Ser Leu Thr Cys Leu Ala Gly Asn Val Ser Ala Cys Asp
 345 350 355

gcc cct att ctc tcc tct tcc agg tcc cta gac tgc agg gag agt ggc 1204
 Ala Pro Ile Leu Ser Ser Arg Ser Leu Asp Cys Arg Glu Ser Gly
 360 365 370

aag aat ggg cct cat gtg tac cag gac ctc ctg ctt agc ctt ggg act 1252
 Lys Asn Gly Pro His Val Tyr Gln Asp Leu Leu Leu Ser Leu Gly Thr
 375 380 385 390

aca aac agc acg ctg ccc cct cca ttt tct ctc caa tct gga atc ctg 1300
 Thr Asn Ser Thr Leu Pro Pro Pro Phe Ser Leu Gln Ser Gly Ile Leu
 395 400 405

aca ttg aac cca gtt gct cag ggt cag ccc att ctt act tcc ctg gga 1348
 Thr Leu Asn Pro Val Ala Gln Gly Gln Pro Ile Leu Thr Ser Leu Gly
 410 415 420

tca aat caa gaa gaa gca tat gtc acc atg tcc agc ttc tac caa aac 1396
 Ser Asn Gln Glu Glu Ala Tyr Val Thr Met Ser Ser Phe Tyr Gln Asn
 425 430 435

cag tgaagtgtaa gaaaccaga ctgaacttac cgtgagcgac aaagatgatt 1449
 Gln

taaaaggga gtctagagtt cctagtctcc ctacagcac agagaagaca aaattagcaa 1509

50 aacccacta cacagtctgc aagattctga aacattgctt tgaccactct tcttgagttc 1569

agtggcactc aacatgagtc aagagcatcc tgcttctacc atgtggattt ggtcacaagg 1629

55 tttaagggtga cccaatgatt cagctattt 1658

<211> 459
 <212> PRT
 <213> Unknown

<400> 2
 Met Thr Ile Leu Gly Thr Thr Phe Gly Met Val Phe Ser Leu Leu Gln -5
 -15
 -20
 Val Val Ser Gly Glu Ser Gly Tyr Ala Gln Asn Gly Asp Leu Glu Asp 10
 -1 1 5
 Ala Glu Leu Asp Asp Tyr Ser Phe Ser Cys Tyr Ser Gln Leu Glu Val 25
 15 20
 Asn Gly Ser Gln His Ser Leu Thr Cys Ala Phe Glu Asp Pro Asp Val 40
 30 35
 Asn Thr Thr Asn Leu Glu Phe Glu Ile Cys Gly Ala Leu Val Glu Val 60
 45 50 55
 Lys Cys Leu Asn Phe Arg Lys Leu Gln Glu Ile Tyr Phe Ile Glu Thr 75
 65 70
 Lys Lys Phe Leu Leu Ile Gly Lys Ser Asn Ile Cys Val Lys Val Gly 90
 80 85
 Glu Lys Ser Leu Thr Cys Lys Lys Ile Asp Leu Thr Thr Ile Val Lys 105
 95 100
 Pro Glu Ala Pro Phe Asp Leu Ser Val Ile Tyr Arg Glu Gly Ala Asn 120
 110 115
 Asp Phe Val Val Thr Phe Asn Thr Ser His Leu Gln Lys Lys Tyr Val 140
 125 130 135
 Lys Val Leu Met His Asp Val Ala Tyr Arg Gln Glu Lys Asp Glu Asn 155
 145 150
 Lys Trp Thr His Val Asn Leu Ser Ser Thr Lys Leu Thr Leu Leu Gln 170
 160 165
 Arg Lys Leu Gln Pro Ala Ala Met Tyr Glu Ile Lys Val Arg Ser Ile 185
 175 180
 Pro Asp His Tyr Phe Lys Gly Phe Trp Ser Glu Trp Ser Pro Ser Tyr 200
 190 195
 Tyr Phe Arg Thr Pro Glu Ile Asn Asn Ser Ser Gly Glu Met Asp Pro 220
 205 210 215
 Ile Leu Leu Thr Ile Ser Ile Leu Ser Phe Phe Ser Val Ala Leu Leu 235
 225 230
 Val Ile Leu Ala Cys Val Leu Trp Lys Lys Arg Ile Lys Pro Ile Val 250
 240 245
 Trp Pro Ser Leu Pro Asp His Lys Lys Thr Leu Glu His Leu Cys Lys 265
 255 260

Lys Pro Arg Lys Asn Leu Asn Val Ser Phe Asn Pro Glu Ser Phe Leu
 270 275 280
 Asp Cys Gln Ile His Arg Val Asp Asp Ile Gln Ala Arg Asp Glu Val
 285 290 295 300
 Glu Gly Phe Leu Gln Asp Thr Phe Pro Gln Gln Leu Glu Glu Ser Glu
 305 310 315
 Lys Gln Arg Leu Gly Gly Asp Val Gln Ser Pro Asn Cys Pro Ser Glu
 320 325 330
 Asp Val Val Val Thr Pro Glu Ser Phe Gly Arg Asp Ser Ser Leu Thr
 335 340 345
 Cys Leu Ala Gly Asn Val Ser Ala Cys Asp Ala Pro Ile Leu Ser Ser
 350 355 360
 Ser Arg Ser Leu Asp Cys Arg Glu Ser Gly Lys Asn Gly Pro His Val
 365 370 375 380
 Tyr Gln Asp Leu Leu Leu Ser Leu Gly Thr Thr Asn Ser Thr Leu Pro
 385 390 395
 Pro Pro Phe Ser Leu Gln Ser Gly Ile Leu Thr Leu Asn Pro Val Ala
 400 405 410
 Gln Gly Gln Pro Ile Leu Thr Ser Leu Gly Ser Asn Gln Glu Glu Ala
 415 420 425
 Tyr Val Thr Met Ser Ser Phe Tyr Gln Asn Gln
 430 435

<210> 3
 <211> 1557
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism: primate; surmised
 human

<220>
 <221> CDS
 <222> (13)..(1125)

<220>
 <221> mat_peptide
 <222> (79)..(1125)

<400> 3
 cggcaccgagg gc atg ggg cgg ctg gtt ctg ctg tgg gga gct gcc gtc ttt 51
 Met Gly Arg Leu Val Leu Leu Trp Gly Ala Ala Val Phe
 -20 -15 -10

ctg ctg gga ggc tgg atg gct ttg ggg caa gga gga gca gca gaa gga 99

Leu Leu Gly Gly Trp Met Ala Leu Gly Gln Gly Gly Ala Ala Glu Gly
 -5 -1 1 5
 gta cag att cag atc atc tac ttc aat tta gaa acc gtg cag gtg aca 147
 Val Gln Ile Gln Ile Ile Tyr Phe Asn Leu Glu Thr Val Gln Val Thr
 10 15 20
 tgg aat gcc agc aaa tac tcc agg acc aac ctg act ttc cac tac aga 195
 Trp Asn Ala Ser Lys Tyr Ser Arg Thr Asn Leu Thr Phe His Tyr Arg
 25 30 35
 ttc aac ggt gat gag gcc tat gac cag tgc acc aac tac ctt ctc cag 243
 Phe Asn Gly Asp Glu Ala Tyr Asp Gln Cys Thr Asn Tyr Leu Leu Gln
 40 45 50 55
 gaa ggt cac act tcg ggg tgc ctc cta gac gca gag cag cga gac gac 291
 Glu Gly His Thr Ser Gly Cys Leu Leu Asp Ala Glu Gln Arg Asp Asp
 60 65 70
 att ctc tat ttc tcc atc agg aat ggg acg cac ccc gtt ttc acc gca 339
 Ile Leu Tyr Phe Ser Ile Arg Asn Gly Thr His Pro Val Phe Thr Ala
 75 80 85
 agt cgc tgg atg gtt tat tac ctg aaa ccc agt tcc ccg aag cac gtg 387
 Ser Arg Trp Met Val Tyr Tyr Leu Lys Pro Ser Ser Pro Lys His Val
 90 95 100
 aga ttt tcg tgg cat cag gat gca gtg acg gtg acg tgt tct gac ctg 435
 Arg Phe Ser Trp His Gln Asp Ala Val Thr Val Thr Cys Ser Asp Leu
 105 110 115
 tcc tac ggg gat ctc ctc tat gag gtt cag tac cgg agc ccc ttc gac 483
 Ser Tyr Gly Asp Leu Leu Tyr Glu Val Gln Tyr Arg Ser Pro Phe Asp
 120 125 130 135
 acc gag tgg cag tcc aaa cag gaa aat acc tgc aac gtc acc ata gaa 531
 Thr Glu Trp Gln Ser Lys Gln Glu Asn Thr Cys Asn Val Thr Ile Glu
 140 145 150
 ggc ttg gat gcc gag aag tgt tac tct ttc tgg gtc agg gtg aag gct 579
 Gly Leu Asp Ala Glu Lys Cys Tyr Ser Phe Trp Val Arg Val Lys Ala
 155 160 165
 atg gag gat gta tat ggg cca gac aca tac cca agc gac tgg tca gag 627
 Met Glu Asp Val Tyr Gly Pro Asp Thr Tyr Pro Ser Asp Trp Ser Glu
 170 175 180
 gtg aca tgc tgg cag aga ggc gag att cgg gat gcc tgt gca gag aca 675
 Val Thr Cys Trp Gln Arg Gly Glu Ile Arg Asp Ala Cys Ala Glu Thr
 185 190 195
 cca acg cct ccc aaa cca aag ctg tcc aaa ttt att tta att tcc agc 723
 Pro Thr Pro Pro Lys Pro Lys Leu Ser Lys Phe Ile Leu Ile Ser Ser
 200 205 210 215
 ctg gcc atc ctt ctg atg gtg tct ctc ctc ctt ctg tct tta tgg aaa 771
 Leu Ala Ile Leu Leu Met Val Ser Leu Leu Leu Leu Ser Leu Trp Lys
 220 225 230

tta tgg aga gtg aag aag ttt ctc att ccc agc gtg cca gac ccg aaa 819
 Leu Trp Arg Val Lys Lys Phe Leu Ile Pro Ser Val Pro Asp Pro Lys
 235 240 245
 tcc atc ttc ccc ggg ctc ttt gag ata cac caa ggg aac ttc cag gag 867
 Ser Ile Phe Pro Gly Leu Phe Glu Ile His Gln Gly Asn Phe Gln Glu
 250 255 260
 tgg atc aca gac acc cag aac gtg gcc cac ctc cac aag atg gca ggt 915
 Trp Ile Thr Asp Thr Gln Asn Val Ala His Leu His Lys Met Ala Gly
 265 270 275
 gca gag caa gaa agt ggc ccc gag gag ccc ctg gta gtc cag ttg gcc 963
 Ala Glu Gln Glu Ser Gly Pro Glu Glu Pro Leu Val Val Gln Leu Ala
 280 285 290 295
 aag act gaa gcc gag tct ccc agg atg ctg gac cca cag acc gag gag 1011
 Lys Thr Glu Ala Glu Ser Pro Arg Met Leu Asp Pro Gln Thr Glu Glu
 300 305 310
 aaa gag gcc tct ggg gga tcc ctc cag ctt ccc cac cag ccc ctc caa 1059
 Lys Glu Ala Ser Gly Gly Ser Leu Gln Leu Pro His Gln Pro Leu Gln
 315 320 325
 ggc ggt gat gtg gtc aca atc ggg ggc ttc acc ttt gtg atg aat gac 1107
 Gly Gly Asp Val Val Thr Ile Gly Gly Phe Thr Phe Val Met Asn Asp
 330 335 340
 cgc tcc tac gtg gcg ttg tgatggacac accactgtca aagtcaacgt 1155
 Arg Ser Tyr Val Ala Leu
 345
 caggatccac gttgacattt aaagacagag gggactgtcc cggggactcc acaccacat 1215
 ggatgggaag tctccacgcc aatgatggta ggactaggag actctgaaga cccagcctca 1275
 ccgcctaagt cgccactgc cctgctaact ttccccaca tgagtctctg tgttcaaagg 1335
 cttgatggca gatgggagcc aattgctcca ggagatttac tcccagttcc ttttcgtgcc 1395
 tgaacgttgt cacataaacc ccaaggcagc acgtccaaaa tgctgtaaaa ccatcttccc 1455
 actctgtgag tcccagttc cgtccatgta cctgttccat agcattggat tctcggagga 1515
 ttttttgtct gttttgagac tccaaaccac ctctaccct ac 1557

<210> 4
 <211> 371
 <212> PRT
 <213> Unknown

<400> 4
 Met Gly Arg Leu Val Leu Leu Trp Gly Ala Ala Val Phe Leu Leu Gly
 -20 -15 -10

Gly Trp Met Ala Leu Gly Gln Gly Gly Ala Ala Glu Gly Val Gln Ile

[illegible]

Ser Gly Gly Ser Leu Gln Leu Pro His Gln Pro Leu Gln Gly Gly Asp
 315 320 325 330
 Val Val Thr Ile Gly Gly Phe Thr Phe Val Met Asn Asp Arg Ser Tyr
 335 340 345
 Val Ala Leu

<210> 5
 <211> 480
 <212> DNA
 <213> Unknown

<220>
 <223> Description of Unknown Organism: primate; surmised
 human

<220>
 <221> CDS
 <222> (1)..(477)

<220>
 <221> mat_peptide
 <222> (85)..(477)

<400> 5
 atg ttc cct ttt gcc tta cta tat gtt ctg tca gtt tct ttc agg aaa 48
 Met Phe Pro Phe Ala Leu Leu Tyr Val Leu Ser Val Ser Phe Arg Lys
 -25 -20 -15
 atc ttc atc tta caa ctt gta ggg ctg gtg tta act tac gac ttc act 96
 Ile Phe Ile Leu Gln Leu Val Gly Leu Val Leu Thr Tyr Asp Phe Thr
 -10 -5 -1 1
 aac tgt gac ttt gag aag att aaa gca gcc tat ctc agt act att tct 144
 Asn Cys Asp Phe Glu Lys Ile Lys Ala Ala Tyr Leu Ser Thr Ile Ser
 5 10 15 20
 aaa gac ctg att aca tat atg agt ggg acc aaa agt acc gag ttc aac 192
 Lys Asp Leu Ile Thr Tyr Met Ser Gly Thr Lys Ser Thr Glu Phe Asn
 25 30 35
 aac acc gtc tct tgt agc aat cgg cca cat tgc ctt act gaa atc cag 240
 Asn Thr Val Ser Cys Ser Asn Arg Pro His Cys Leu Thr Glu Ile Gln
 40 45 50
 agc cta acc ttc aat ccc acc gcc ggc tgc gcg tgc ctc gcc aaa gaa 288
 Ser Leu Thr Phe Asn Pro Thr Ala Gly Cys Ala Ser Leu Ala Lys Glu
 55 60 65
 atg ttc gcc atg aaa act aag gct gcc tta gct atc tgg tgc cca ggc 336
 Met Phe Ala Met Lys Thr Lys Ala Ala Leu Ala Ile Trp Cys Pro Gly
 70 75 80
 tat tcg gaa act cag ata aat gct act cag gca atg aag aag agg aga 384

Tyr Ser Glu Thr Gln Ile Asn Ala Thr Gln Ala Met Lys Lys Arg Arg 100
 85 90 95
 aaa agg aaa gtc aca acc aat aaa tgt ctg gaa caa gtg tca caa tta 432
 Lys Arg Lys Val Thr Thr Asn Lys Cys Leu Glu Gln Val Ser Gln Leu 115
 105 110
 caa gga ttg tgg cgt cgc ttc aat cga cct tta ctg aaa caa cag taa 480
 Gln Gly Leu Trp Arg Arg Phe Asn Arg Pro Leu Leu Lys Gln Gln 130
 120 125

<210> 6
 <211> 159
 <212> PRT
 <213> Unknown

<400> 6
 Met Phe Pro Phe Ala Leu Leu Tyr Val Leu Ser Val Ser Phe Arg Lys -15
 -25 -20

Ile Phe Ile Leu Gln Leu Val Gly Leu Val Leu Thr Tyr Asp Phe Thr
 -10 -5 -1 1

Asn Cys Asp Phe Glu Lys Ile Lys Ala Ala Tyr Leu Ser Thr Ile Ser 20
 5 10 15

Lys Asp Leu Ile Thr Tyr Met Ser Gly Thr Lys Ser Thr Glu Phe Asn 35
 25 30

Asn Thr Val Ser Cys Ser Asn Arg Pro His Cys Leu Thr Glu Ile Gln 50
 40 45

Ser Leu Thr Phe Asn Pro Thr Ala Gly Cys Ala Ser Leu Ala Lys Glu 65
 55 60

Met Phe Ala Met Lys Thr Lys Ala Ala Leu Ala Ile Trp Cys Pro Gly 80
 70 75

Tyr Ser Glu Thr Gln Ile Asn Ala Thr Gln Ala Met Lys Lys Arg Arg 100
 85 90 95

Lys Arg Lys Val Thr Thr Asn Lys Cys Leu Glu Gln Val Ser Gln Leu 115
 105 110

Gln Gly Leu Trp Arg Arg Phe Asn Arg Pro Leu Leu Lys Gln Gln 130
 120 125